

# Cl-IB-MECA [2-Chloro-*N*<sup>6</sup>-(3-iodobenzyl)adenosine-5'-*N*-methylcarboxamide] Reduces Ischemia/Reperfusion Injury in Mice by Activating the A<sub>3</sub> Adenosine Receptor<sup>S</sup>

Zhi-Dong Ge, Jason N. Peart, Laura M. Kreckler, Tina C. Wan, Marlene A. Jacobson, Garrett J. Gross, and John A. Auchampach

Department of Pharmacology and Toxicology and the Cardiovascular Center (Z.-D.G., J.N.P., L.M.K., T.C.W. G.J.G., J.A.A.), Medical College of Wisconsin, Milwaukee, Wisconsin; and Department of Pharmacology, Merck Research Laboratories, West Point, Pennsylvania (M.A.J.)

Received July 21, 2006; accepted September 18, 2006

## ABSTRACT

We used pharmacological agents and genetic methods to determine whether the potent A<sub>3</sub> adenosine receptor (AR) agonist 2-chloro-*N*<sup>6</sup>-(3-iodobenzyl)adenosine-5'-*N*-methylcarboxamide (Cl-IB-MECA) protects against myocardial ischemia/reperfusion injury in mice via the A<sub>3</sub>AR or via interactions with other AR subtypes. Pretreating wild-type (WT) mice with Cl-IB-MECA reduced myocardial infarct size induced by 30 min of coronary occlusion and 24 h of reperfusion at doses (30 and 100  $\mu$ g/kg) that concomitantly reduced blood pressure and stimulated systemic histamine release. The A<sub>3</sub>AR-selective antagonist MRS 1523 [3-propyl-6-ethyl-5[(ethylthio)carbonyl]-2-phenyl-4-propyl-3-pyridine-carboxylate], but not the A<sub>2A</sub>AR antagonist ZM 241385 [4-{2-[7-amino-2-(2-furyl)[1,2,4]triazolo-[2,3-a][1,3,5]triazin-5-ylamino]ethyl}phenol], blocked the

reduction in infarct size provided by Cl-IB-MECA, suggesting a mechanism involving the A<sub>3</sub>AR. To further examine the selectivity of Cl-IB-MECA, we assessed its cardioprotective effectiveness in A<sub>3</sub>AR gene "knock-out" (A<sub>3</sub>KO) mice. Cl-IB-MECA did not reduce myocardial infarct size in A<sub>3</sub>KO mice *in vivo* and did not protect isolated perfused hearts obtained from A<sub>3</sub>KO mice from injury induced by global ischemia and reperfusion. Additional studies using WT mice treated with compound 48/80 [condensation product of *p*-methoxyphenethyl methylamine with formaldehyde] to deplete mast cell contents excluded the possibility that Cl-IB-MECA was cardioprotective by releasing mediators from mast cells. These data demonstrate that Cl-IB-MECA protects against myocardial ischemia/reperfusion injury in mice principally by activating the A<sub>3</sub>AR.

Several different A<sub>3</sub> adenosine receptor (AR) agonists, including the *N*<sup>6</sup>-benzyl adenosine-5'-*N*-methylcarboxamide derivatives IB-MECA, Cl-IB-MECA, and CB-MECA, have been shown to be effective at protecting against myocardial ischemia/reperfusion injury in animal models of infarction and myocardial stunning (Auchampach et al., 1997b, 2003; Tracey et al., 1997, 1998, 2003; Jordan et al., 1999; Thourani

et al., 1999a,b; Kodani et al., 2001; Takano et al., 2001). However, it remains uncertain whether these agents are effective by activating the A<sub>3</sub>AR or by nonspecific interactions with other AR subtypes. This issue has been difficult to address, because useful A<sub>3</sub>AR antagonists have only recently been developed.

The goal of this investigation was to test the cardioprotective effectiveness of Cl-IB-MECA in an *in vivo* mouse model of infarction and in an isolated mouse heart model of global ischemia and reperfusion. A second goal of this investigation was to establish definitively whether Cl-IB-MECA exerts cardioprotection by activating A<sub>3</sub>ARs. Our experimental approach involved the use of the rodent A<sub>3</sub>AR antagonist MRS 1523 (Li et al., 1998), the potent A<sub>2A</sub>AR antagonist ZM

This work was supported by National Institutes of Health Grants R01 HL60051, R01 HL07707, and T32 HL73643 and by American Heart Association Research Fellowships 0320019Z and 0315274Z.

Article, publication date, and citation information can be found at <http://jpet.aspetjournals.org>.

doi:10.1124/jpet.106.111351.

<sup>S</sup> The online version of this article (available at <http://jpet.aspetjournals.org>) contains supplemental material.

**ABBREVIATIONS:** AR, adenosine receptor; CB-MECA, *N*<sup>6</sup>-(3-chlorobenzyl)adenosine-5'-*N*-methylcarboxamide; Cl-IB-MECA, 2-chloro-*N*<sup>6</sup>-(3-iodobenzyl)adenosine-5'-*N*-methylcarboxamide; [<sup>125</sup>I]I-AB-MECA; *N*<sup>6</sup>-(4-amino-3-[<sup>125</sup>I]iodobenzyl)adenosine-5'-*N*-methylcarboxamide; LAD, left anterior descending; LV, left ventricle; KO, knock-out; PE, polyethylene; WT, wild-type; HEK, human embryonic kidney; MRS 1523, 3-propyl-6-ethyl-5[(ethylthio)carbonyl]-2-phenyl-4-propyl-3-pyridine-carboxylate; ZM 241385, 4-{2-[7-amino-2-(2-furyl)[1,2,4]triazolo-[2,3-a][1,3,5]triazin-5-ylamino]ethyl}phenol; compound 48/80, condensation product of *p*-methoxyphenethyl methylamine with formaldehyde; SKF-91488, 4-(*N,N*-dimethylamino)butylisothiourea; dP/dt, change in pressure over time; CGS 21680, 2-[*p*-(2-carboxyethyl)phenylethylamino]-5'-*N*-ethylcarboxamidoadenosine.

241385, and mice with genetic deletion of the A<sub>3</sub>AR gene (A<sub>3</sub>KO mice; Salvatore et al., 2000). We examined whether cardioprotection provided by Cl-IB-MECA is sensitive to blockade by ZM 241385, because the possibility has been raised that N<sup>6</sup>-benzyladenosine-5'-N-methylcarboxamide A<sub>3</sub>AR agonists may be beneficial in animal models of tissue injury via interactions with the A<sub>2A</sub>AR rather than the A<sub>3</sub>AR (Murphree et al., 2002; Lappas et al., 2005). Finally, because A<sub>3</sub>AR agonists induce mast cell degranulation in rodents (Linden, 1994; Hannon et al., 1995; Van Schaik et al., 1996), we also examined the cardioprotective profile of Cl-IB-MECA in mice that had been depleted of mast cell contents by chronic treatment with compound 48/80.

## Materials and Methods

### Materials

Cell culture reagents G418, pcDNA3.1, and Lipofectamine were purchased from Invitrogen (Carlsbad, CA). ZM 241385 was from Tocris Cookson, Inc. (Ellisville, MO), adenosine deaminase was from Roche Applied Science (Indianapolis, IN), and all remaining drugs and reagents were purchased from Sigma-Aldrich (St. Louis, MO). N<sup>6</sup>-(4-Amino-3-[<sup>125</sup>I]iodobenzyl)adenosine-5'-N-methylcarboxamide ([<sup>125</sup>I]I-AB-MECA) was synthesized and purified by HPLC, as described previously (Olah et al., 1994; Auchampach et al., 1997a). cAMP and histamine radioimmunoassay kits were obtained from GE Healthcare (Piscataway, NJ) and Immunotech (Marseille, France), respectively.

### Animals

All experiments were performed with 10- to 14-week-old male mice (weighing ~25–30 g). Wild-type (WT) FVB/N and C57BL/6 mice were purchased from Taconic Farms Inc. (Germantown, NY). A<sub>3</sub>KO mice were generated by embryonic stem cell targeting and genotyped by Southern blotting, as described previously (Salvatore et al., 2000). A<sub>3</sub>KO mice used had been transferred to the C57BL/6 genetic background by backcrossing greater than 12 generations. Transgenic mice cardiac-specifically overexpressing the A<sub>3</sub>AR (A<sub>3</sub>tg.1) were generated on the FVB/N genetic background, as described previously (Black et al., 2002). All animals in the study received humane care in accordance with the guidelines established by the Medical College of Wisconsin, which conform to the *Guide for the Care and Use of Laboratory Animals* published by the US National Institutes of Health (NIH Publication 85-23, revised 1996).

### Radioligand Binding Assays and cAMP Accumulation Assays

Competition radioligand binding assays using [<sup>125</sup>I]I-AB-MECA were conducted with membranes prepared from HEK 293 cells expressing recombinant mouse A<sub>1</sub> or A<sub>3</sub>ARs, and cAMP accumulation assays were performed with HEK 293 cells transfected with mouse A<sub>2A</sub> or A<sub>2B</sub>ARs, as described previously (Auchampach et al., 1997a; Takano et al., 2001; Kreckler et al., 2006).

### In Vivo Mouse Model of Infarction

**Surgical Preparation.** Mice were anesthetized with sodium pentobarbital (100 mg/kg i.p.) and placed on a warm heating pad to maintain body temperature at 37 ± 0.3°C. A polyethylene (PE)-60 tube was inserted into the trachea and connected to a mouse ventilator (model 845; Hugo Sachs Elektronic, Hugstetten, Germany). The mice were respiration (tidal volume = 225 µl; rate ~100 strokes/min) with room air supplemented with 100% oxygen to maintain blood gases within normal physiological limits. The electrocardiogram (ECG) was obtained with needle electrodes using the limb lead II configuration and recorded continuously using a Powerlab data acquisition system (ADIInstruments, Colorado Springs, CO).

A left thoracotomy was performed at the fourth intercostal space to expose the heart, the pericardium was removed, and an 8.0 nylon suture was passed below the left anterior descending (LAD) coronary artery 1 to 3 mm from the tip of the left atrium with the aid of a dissecting microscope. Myocardial ischemia was induced by tying the suture over a piece of wetted gauze, and reperfusion was initiated by loosening the suture. Successful performance of coronary occlusion and reperfusion was verified by visual inspection (i.e., by the development of a pale color in the distal myocardium upon occlusion and the return of a bright red color due to hyperemia after release) and by observing changes in the ECG (i.e., widening of the QRS complex and ST segment changes). The chest wall was then closed with a 7-0 polypropylene suture with one layer through the chest wall and muscle and a second layer through the skin and subcutaneous tissue. Subsequently, the mouse was removed from the ventilator and kept in a warm chamber supplied with 100% oxygen. The endotracheal tube was removed after approximately 60 min when the mice began to recover their righting reflex and respiratory rate was ~140 breaths/min.

**Experimental Protocol.** After a 30-min stabilization period, all mice were subjected to 30 min of coronary occlusion and 24 h of reperfusion. Cl-IB-MECA or equivalent vehicle (0.2 ml of 50% dimethyl sulfoxide in normal saline) was administered as i.v. boluses (tail vein injection) at the doses indicated beginning 10 min before the coronary occlusion. In studies using AR antagonists, the drugs were administered by i.v. injection 15 min before the administration of Cl-IB-MECA. Heart rate was monitored at baseline and during the 30-min occlusion period from the ECG recording.

**Measurement of Ischemic Area and Infarct Size.** After 24 h of reperfusion, the mice were administered heparin (1 U/g i.p.) and anesthetized with pentobarbital. After anesthesia, a thoracotomy was performed and the ascending aorta was cannulated with PE-10 tubing. To delineate the ischemic area at risk, the LAD was reoccluded and a 5% solution of phthalo blue dye in normal saline was injected into the aortic root. The heart was excised, washed with phosphate-buffered saline, embedded in agarose, and frozen at -20°C. To delineate infarcted tissue, the left ventricle was sliced into five to six transverse pieces using a microtome, incubated in a 1% solution of triphenyltetrazolium chloride for 10 min at 37°C to stain viable tissue red, and fixed in 10% formaldehyde overnight. The slices were weighed and photographed from both sides using a dissecting microscope mounted with a SPOT Insight digital camera (Diagnostic Instruments, Inc., Sterling Heights, MI). The left ventricular, ischemic, and infarcted areas were measured by digital planimetry. Infarct size is presented as a percentage of the ischemic risk region.


**Measurement of Systemic Blood Pressure and Plasma Histamine Levels.** Parallel experiments were performed on two separate groups of mice to determine the effect of Cl-IB-MECA on systemic blood pressure and plasma histamine levels. For blood pressure measurements, the mice were anesthetized with pentobarbital and respiration as described above. A saline-filled cannula (stretched PE-10 tubing) was inserted into the left femoral artery and connected to a pressure transducer (ADIInstruments model MLT 0699). Blood pressure was recorded continuously by a PowerLab data acquisition system (ADIInstruments).

For measurement of plasma histamine levels, the mice were anesthetized, connected to the respirator, and subjected to a thoracotomy to expose the heart. Blood samples (0.15 ml) were collected by cardiac puncture into EDTA-coated syringes containing the histamine N-methyltransferase inhibitor SKF-91488 (final concentration = 20 µM) at baseline and at 15, 30, and 45 min after the administration of Cl-IB-MECA. The blood samples were centrifuged (1000g for 10 min at 4°C), and the plasma was analyzed for histamine content by radioimmunoassay.

### Langendorff-Perfused Mouse Heart Model of Global Ischemia and Reperfusion Experimental Setup.

Mice were anesthetized with pentobarbital. As soon as deep anesthesia was achieved, the hearts were excised quickly and arrested in ice-cold perfusion buffer. The hearts were cannulated (20 gauge stainless steel) via the aorta and perfused by the Langendorff method using Krebs-Henseleit buffer containing 118 mM NaCl, 25 mM NaHCO<sub>3</sub>, 4.7 mM KCl, 1.2 mM MgCl<sub>2</sub>, 2.5 mM CaCl<sub>2</sub>, 1.2 mM KH<sub>2</sub>PO<sub>4</sub>, 0.5 mM EDTA, and 11 mM glucose. The buffer was equilibrated with 95% O<sub>2</sub>/5% CO<sub>2</sub> at 37°C to maintain the pH at 7.4. The left ventricle was ventilated with a PE drain, and a fluid-filled balloon was inserted into the ventricle via the mitral valve. The balloon was connected to a pressure transducer for continuous measurement of left ventricular pressure. The hearts were immersed in perfusate maintained at 37°C, and the balloon was inflated to a diastolic pressure of ~5 to 10 mm Hg. Coronary flow was monitored by an in-line flow probe connected to a flowmeter (Model T206; Transonics Systems Inc., Ithaca, NY). The left ventricular pressure signal was acquired by a PowerLab data acquisition system (ADInstruments) and processed to yield heart rate and left ventricular dP/dt.

**Protocol.** Isolated hearts were perfused for 20 min to allow for stabilization and then perfused for an additional 15 min while pacing at 420 beats/min (ventricular pacing, 2-ms square waves, 20% above threshold). Baseline measurements of function were acquired immediately before subjecting the hearts to 20 min of no-flow global ischemia and 45 min of reperfusion. To examine the effect of Cl-IB-MECA on functional recovery, hearts were perfused with buffer dP/dt.



containing 100 nM Cl-IB-MECA for 10 min before ischemia and throughout the reperfusion period.

### Statistical Analysis

$K_i$  and EC<sub>50</sub> values are reported as geometrical means with corresponding 95% confidence intervals. All other data are presented as arithmetic means  $\pm$  S.E.M. Hemodynamic variables and plasma histamine concentrations were analyzed by two-way repeated measures ANOVA (time and treatment) to determine whether there was a main effect of time, a main effect of treatment, or a time-treatment interaction. If global tests showed a main effect or interaction, post hoc tests were performed using unpaired or paired analyses, as appropriate. Infarct size, ischemic area at risk size, and left ventricular functional recoveries in the isolated heart studies were compared by one-way ANOVA followed by a Student's *t* test with the Bonferroni's correction or an unpaired *t* test, as appropriate.

## Results


**Selectivity of Cl-IB-MECA for Recombinant Mouse ARs.** Preliminary assays were conducted with transfected HEK 293 cells to determine the affinity and selectivity of Cl-IB-MECA and its parent compound IB-MECA for recombinant mouse ARs. Competition radioligand binding assays were performed to compare the binding affinity of the ligands for A<sub>1</sub> and A<sub>3</sub>ARs. Because high-affinity agonist binding to A<sub>2A</sub>ARs can not be accurately measured in heterologous ex-

pression systems due to poor receptor coupling (Murphree et al., 2002) and because agonist radioligands are not available for the A<sub>2B</sub>AR, we determined the potency of Cl-IB-MECA and IB-MECA to stimulate cAMP production in HEK 293 cells expressing mouse A<sub>2A</sub> or A<sub>2B</sub>ARs. On the basis of radioligand binding analysis, Cl-IB-MECA was determined to be 219-fold selective at binding to the high-affinity form of the mouse A<sub>3</sub>AR versus the high-affinity form of the mouse A<sub>1</sub>AR (Fig. 1). The dissociation constants were calculated to be 35 nM (95% confidence intervals = 33 and 36 nM) for the mouse A<sub>1</sub>AR and 0.18 nM (0.16 and 0.19 nM) for the mouse A<sub>3</sub>AR. In comparison, IB-MECA was 68-fold more potent at binding to the mouse A<sub>3</sub>AR versus the mouse A<sub>1</sub>AR [5.9 nM (4.4 and 7.9 nM) versus 0.087 nM (0.078 and 0.098 nM), respectively]. In cAMP assays (Fig. 1), Cl-IB-MECA and IB-MECA displayed low potency at stimulating cAMP production in HEK 293 cells expressing either A<sub>2A</sub> or A<sub>2B</sub>ARs. In fact, accurate EC<sub>50</sub> values could not be calculated for these ligands due to failure to reach maximal responses at their solubility limit of 10  $\mu$ M. It is reasonable to estimate that Cl-IB-MECA was at least 1000-fold less potent at stimulating cAMP production in A<sub>2A</sub>AR-expressing cells compared with the potent A<sub>2A</sub>AR agonist CGS 21680. In A<sub>2B</sub>AR-expressing cells, Cl-IB-MECA was considerably less potent at stimulating cAMP (NECA) production compared with adenosine-5'-N-ethylcarboxamide, the most potent A<sub>2B</sub>AR agonist currently known.

**Dose-Dependent Reduction in Infarct Size by Cl-IB-MECA.** We initially conducted a dose-response study (10–100  $\mu$ g/kg) with Cl-IB-MECA examining its ability to reduce infarct size in WT FVB/N mice in response to 30 min of LAD occlusion and 24 h of reperfusion. In separate groups of mice, we assessed the effect of Cl-IB-MECA on systemic blood pressure and plasma histamine concentrations. The data are presented in Fig. 2. In WT FVB/N mice, administration of Cl-IB-MECA produced a dose-dependent reduction in infarct size that was significantly different from vehicle at 30  $\mu$ g/kg. At a dose of 100  $\mu$ g/kg, the magnitude of the reduction in infarct size provided by Cl-IB-MECA was 36% compared with vehicle-treated mice (30  $\pm$  3% of the ischemic area at risk versus 47  $\pm$  2%, respectively). Treatment with Cl-IB-MECA had no effect on heart rate during the 30-min ischemic period (Table 1).

In parallel experiments, we observed that Cl-IB-MECA also reduced blood pressure and significantly increased plasma histamine levels (Fig. 2). The decrease in blood pressure was transient peaking at 10 min after Cl-IB-MECA administration (15% reduction by 100  $\mu$ g/kg Cl-IB-MECA) and returned to normal within 30 min. The increase in plasma histamine concentration was most prominent at a dose of 100  $\mu$ g/kg (from 151  $\pm$  16 to 1763  $\pm$  298 nM 15 min after Cl-IB-MECA administration), although a small increase was also detected at a dose of 30  $\mu$ g/kg (from 150  $\pm$  11 to 358  $\pm$  132 nM).

**Cl-IB-MECA Reduced Infarct Size in A<sub>3</sub>AR Transgenic Mice.** We have previously created transgenic mice that cardiac-specifically overexpress the A<sub>3</sub>AR using the  $\alpha$ -myosin heavy chain gene promoter and have shown that these mice demonstrate increased tolerance to ischemia/reperfusion injury (Black et al., 2002; Cross et al., 2002), supporting the theory that the A<sub>3</sub>AR is capable of coupling to protective signaling pathways in cardiomyocytes. In the present study, we examined whether administration of a




**Fig. 2.** A, myocardial infarct size expressed as a percentage of the risk region induced by 30 min of LAD coronary occlusion and 24 h of reperfusion in WT FVB/N mice treated with increasing concentrations of Cl-IB-MECA. Cl-IB-MECA was administered i.v. 10 min before coronary occlusion. Filled circles indicate the mean  $\pm$  S.E.M. Open circles indicate infarct size in individual animals. \*,  $p$   $<$  0.05 versus the vehicle-treated group. Risk region size ranged from 38  $\pm$  2 to 41  $\pm$  2% of the left ventricle and was not significantly different among the treatment groups. B and C, the effect of increasing doses of Cl-IB-MECA on mean arterial blood pressure expressed as a percentage of baseline and plasma histamine levels (in nanomoles) in FVB/N mice. Baseline mean arterial blood pressure was 88  $\pm$  2 mm Hg. All values are the mean  $\pm$  S.E.M. In B and C,  $n$  = 6–8 mice/group. \*,  $p$   $<$  0.05 versus vehicle or time 0.

cardioprotective dose of Cl-IB-MECA provides additional protection in A<sub>3</sub>AR transgenic mice. We used A<sub>3</sub>tg.1 mice containing a single copy of the  $\alpha$ -myosin heavy chain-A<sub>3</sub>AR transgene expressing 12.5  $\pm$  3.2 fmol/mg of the high-affinity G protein-coupled form of the A<sub>3</sub>AR in the heart (Black et al., 2002). As shown in Fig. 3, administration of 100  $\mu$ g/kg Cl-IB-MECA decreased infarct size in A<sub>3</sub>tg.1 mice from 35  $\pm$  4 to 25  $\pm$  3%. Administration of Cl-IB-MECA caused a 10% re-

TABLE 1  
Heart rate during the ischemia/reperfusion experiments

Treatment	Strain	Genotype	Pre-Occ	Ischemia	
				15 Min	30 Min
Vehicle	FVB/N	WT	380 ± 14	412 ± 10	422 ± 10
Cl-IB-MECA 10 $\mu$ g/kg	FVB/N	WT	370 ± 9	416 ± 9	415 ± 12
Cl-IB-MECA 30 $\mu$ g/kg	FVB/N	WT	369 ± 13	397 ± 15	406 ± 11
Cl-IB-MECA 100 $\mu$ g/kg	FVB/N	WT	385 ± 12	413 ± 16	399 ± 12
Vehicle	FVB/N	WT	382 ± 24	395 ± 25	384 ± 26
Vehicle	FVB/N	A <sub>3</sub> tg0.1	380 ± 25	383 ± 20	374 ± 16
Cl-IB-MECA 100 $\mu$ g/kg	FVB/N	A <sub>3</sub> tg0.1	378 ± 17	345 ± 13*	335 ± 18*
ZM 241385 2.0 mg/kg	FVB/N	WT	376 ± 13	409 ± 12	418 ± 13
Cl-IB-MECA 100 $\mu$ g/kg + ZM 0.5 mg/kg	FVB/N	WT	389 ± 14	411 ± 15	420 ± 14
Cl-IB-MECA 100 $\mu$ g/kg + ZM 2.0 mg/kg	FVB/N	WT	383 ± 15	411 ± 14	421 ± 12
MRS 1523 2.0 mg/kg	FVB/N	WT	369 ± 15	408 ± 6	414 ± 12
Cl-IB-MECA 100 $\mu$ g/kg + MRS 2.0 mg/kg	FVB/N	WT	397 ± 13	417 ± 15	418 ± 8
Vehicle	C57BL/6	A <sub>3</sub> AR <sup>+/+</sup>	376 ± 21	402 ± 24	424 ± 20
Vehicle	C57BL/6	A <sub>3</sub> AR <sup>-/-</sup>	384 ± 24	399 ± 23	406 ± 16
Cl-IB-MECA 100 $\mu$ g/kg	C57BL/6	A <sub>3</sub> AR <sup>+/+</sup>	378 ± 13	393 ± 10	406 ± 9
Cl-IB-MECA 100 $\mu$ g/kg	C57BL/6	A <sub>3</sub> AR <sup>-/-</sup>	381 ± 21	397 ± 17	395 ± 15

\*  $P < 0.05$  vs. the vehicle-treated group for each mouse strain by two-way repeated measures ANOVA (time and drug treatment) and Student's *t* test with the Bonferroni's correction.  $n = 7$  to 10/group.




**Fig. 3.** A, myocardial infarct size (mean ± S.E.M.;  $n = 8$  mice/group) expressed as a percentage of the risk region induced by 30 min of LAD coronary occlusion and 24 h of reperfusion in transgenic mice that cardiac-specifically overexpressed the A<sub>3</sub>AR (A<sub>3</sub>ARtg.1) treated 10 min before coronary occlusion (i.v.) with either vehicle or 100  $\mu$ g/kg Cl-IB-MECA. \*,  $p < 0.05$  versus the vehicle-treated group. Risk region size was 38 ± 1 and 40 ± 2% of the left ventricle for the vehicle- and Cl-IB-MECA-treated groups, respectively. B, representative images of heart slices stained with phthalo blue dye and triphenyltetrazolium chloride used to assess infarct size.

duction in heart rate after 30 min of ischemia (Table 1) that was probably due to activation of exogenously expressed A<sub>3</sub>ARs in pacemakers cells (Black et al., 2002). Although Cl-IB-MECA produced additional cardioprotection in A<sub>3</sub>tg.1 mice, there was no difference in infarct size between A<sub>3</sub>tg.1 and WT mice treated with Cl-IB-MECA reported in Fig. 1.

**Cl-IB-MECA Reduces Infarct Size in Vivo via a Specific Interaction with the A<sub>3</sub>AR.** To provide evidence whether or not Cl-IB-MECA effectively reduces infarct size via an A<sub>3</sub>AR-mediated mechanism, we initially assessed the effect of MRS 1523 (Fig. 4). MRS 1523 is a dihydropyridine compound developed by Li et al. (1998) as a rat A<sub>3</sub>AR antagonist ( $K_i = 113 \pm 12$  nM for rat A<sub>3</sub>ARs; 140-fold selectivity). In a previous study (Kreckler et al., 2006), we determined that MRS 1523 binds to mouse A<sub>3</sub>ARs with high affinity ( $K_i = 731$  nM) and with moderate selectivity versus the A<sub>1</sub>AR (11-fold) but excellent selectivity versus A<sub>2A</sub> and A<sub>2B</sub>ARs (at least 1000-fold). As shown in Fig. 4, pretreatment with 2 mg/kg MRS 1523 blocked the reduction in infarct size produced by 100  $\mu$ g/kg Cl-IB-MECA. This dose of MRS 1523 also inhibited the ability of Cl-IB-MECA to increase plasma histamine concentrations and reduce blood pressure (Fig. 4).


We subsequently examined the effect of the A<sub>2A</sub>AR antagonist ZM 241385 on cardioprotection provided by Cl-IB-MECA. Mice were pretreated with 2 mg/kg ZM 241385 15 min before the administration of 100  $\mu$ g/kg Cl-IB-MECA. This dose of ZM 241385 was based on the study of Ohta and Sitkovsky (2001). As shown in Fig. 5, we were surprised to observe that 2 mg/kg ZM 241385 blocked the beneficial effect of Cl-IB-MECA on infarct size. However, we observed that this dose of ZM 241385 also antagonized the ability of Cl-IB-MECA to increase plasma histamine levels and to reduce blood pressure (Fig. 5), suggesting that ZM 241385 at a dose of 2 mg/kg is sufficient to block A<sub>3</sub>ARs. We subsequently tested ZM 241385 at a lower dose of 0.5 mg/kg. This dose only partially antagonized the hemodynamic and histamine-releasing actions of Cl-IB-MECA (Fig. 5) but blocked completely the hypotensive response induced by i.v. administration of 30  $\mu$ g/kg A<sub>2A</sub>AR agonist CGS 21680 (Supplemental Fig. 1). As shown in Fig. 5, treatment with 0.5 mg/kg ZM 241385 failed to block the infarct size-reducing effect of Cl-IB-MECA.

Finally, we examined whether Cl-IB-MECA reduces infarct size in A<sub>3</sub>KO mice. We used A<sub>3</sub>KO mice made congenic on the C57BL/6 genetic background by back-crossing greater than 12 generations to the pure strain. Comparisons were made with WT C57BL/6 mice. Similar to FVB/N mice, pre-treating WT C57BL/6 mice with 100  $\mu$ g/kg Cl-IB-MECA re-




**Fig. 4.** A, myocardial infarct size expressed as a percentage of the risk region induced by 30 min of LAD coronary occlusion and 24 h of reperfusion in WT FVB/N mice treated 10 min before the coronary occlusion with vehicle or 100 µg/kg Cl-IB-MECA in the presence or absence of the A<sub>3</sub>AR antagonist MRS 1523. Mice were given 2 mg/kg MRS 1523 (i.v.) 15 min before the administration of Cl-IB-MECA. Risk region size ranged from 36 ± 3 to 43 ± 2% of the left ventricle and was not significantly different among the treatment groups. B and C, effect of Cl-IB-MECA (100 µg/kg) on mean arterial blood pressure and plasma histamine levels in FVB/N mice pretreated with 2.0 mg/kg MRS 1523. Baseline mean arterial blood pressure was 89 ± 3 mm Hg. All of the data are presented as the mean ± S.E.M. *n* = 9–10 (A) or 6–8 (B and C) mice/group. \*, *p* < 0.05 versus the vehicle-treated group or time 0.

duced infarct size 54% (vehicle = 52 ± 4% of the risk region; Cl-IB-MECA = 24 ± 2%), decreased systemic blood pressure 15% (with no effect on heart rate), and increased plasma histamine levels (Fig. 6; Table 1). The onset of the hypotensive effect of Cl-IB-MECA in C57BL/6 mice was immediate (within 5 min) but more sustained (>45 min) compared with that produced in FVB/N mice. Infarct size induced by 30 min of occlusion and 24 h of reperfusion was similar in magnitude in A<sub>3</sub>KO mice (56 ± 4% in A<sub>3</sub>KO mice) compared with WT



**Fig. 5.** A, myocardial infarct size expressed as a percentage of the risk region induced by 30 min of LAD coronary occlusion and 24 h of reperfusion in WT FVB/N mice treated 10 min before the coronary occlusion with vehicle or 100 µg/kg Cl-IB-MECA in the presence or absence of the A<sub>2A</sub>AR antagonist ZM 241385. Mice were given 0.5 or 2.0 mg/kg ZM 241385 (i.v.) 15 min before the administration of Cl-IB-MECA. Risk region size ranged from 36 ± 3 to 41 ± 2% of the left ventricle and was not significantly different among the treatment groups. B and C, effect of Cl-IB-MECA (100 µg/kg) on mean arterial blood pressure and plasma histamine levels in FVB/N mice pretreated with 0.5 or 2.0 mg/kg ZM 241385. Baseline mean arterial blood pressure was 88 ± 3 mm Hg. All data are presented as the mean ± S.E.M. *n* = 9–10 (A) or 6–8 (B and C) mice/group. \*, *p* < 0.05 versus the vehicle-treated group or time 0.

mice (Fig. 6). However, unlike WT mice, administration of Cl-IB-MECA (100 µg/kg) to A<sub>3</sub>KO mice subjected to ischemia and reperfusion did not reduce infarct size (infarct size as a percentage of the risk region was 56 ± 2%; Fig. 6); similarly, it did not evoke systemic release of histamine (Fig. 6). These results concur with the data obtained with AR antagonists and indicate that Cl-IB-MECA reduces infarct size by an




**Fig. 6.** A and B, myocardial infarct size expressed as a percentage of the risk region in WT C57BL/6 mice and A<sub>3</sub>KO mice subjected to 30 min of coronary occlusion and 24 h of reperfusion. Cl-IB-MECA was administered i.v. 10 min before coronary occlusion. Risk region size ranged from 37 ± 2 to 42 ± 2% of the left ventricle and was not significantly different among the treatment groups. C to F, mean arterial blood pressure and plasma histamine levels in WT (C and E) and A<sub>3</sub>KO (D and F) mice after administration of 100 µg/kg Cl-IB-MECA. MRS 1523 (2 mg/kg) or ZM 241385 (2 mg/kg) were administered to WT or A<sub>3</sub>KO mice 15 min before the administration of Cl-IB-MECA in some of the hemodynamic studies. Baseline mean arterial blood pressure was 86 ± 4 mm Hg in WT mice and 88 ± 2 mm Hg in A<sub>3</sub>KO mice. All data are presented as the mean ± S.E.M. *n* = 9–12 (A and B) or 6–8 (C and F) mice/group. \*, *p* < 0.05 versus the vehicle-treated group or time 0.

A<sub>3</sub>AR-mediated mechanism. It is interesting that we observed that Cl-IB-MECA continued to reduce blood pressure in A<sub>3</sub>KO mice, although the kinetics of the response was different. Rather than a rapid but sustained hypotensive effect, Cl-IB-MECA produced a delayed reduction in blood pressure that became apparent 15 min after drug administration (Fig. 6). Cl-IB-MECA produced a similar hypotensive profile in WT C57BL/6 mice pretreated with 2 mg/kg A<sub>3</sub>AR antagonist MRS 1523, as well as in A<sub>3</sub>KO mice pretreated with a high dose of ZM 241385 (2 mg/kg) that presumably blocks all four AR subtypes (Fig. 6).

**Cl-IB-MECA Protects Isolated Mouse Hearts from Global Ischemia/Reperfusion Injury via the A<sub>3</sub>AR.** It has previously been reported that Cl-IB-MECA improves reperfusion contractile function of isolated mouse hearts subjected to global ischemia/reperfusion injury (Harrison et al., 2002; Peart et al., 2002). In this study, we examined whether the protective effect of Cl-IB-MECA in isolated hearts involves a specific interaction with the A<sub>3</sub>AR. Langendorff

perfused hearts from WT or A<sub>3</sub>KO mice were subjected to 20 min of normothermic no-flow global ischemia and 45 min of reperfusion; the hearts were treated throughout the experiment with either vehicle or 100 nM Cl-IB-MECA. Postischemic recovery of left ventricular function was assessed via a fluid-filled balloon placed in the left ventricle. As shown in Fig. 7, indices of left ventricular contractile function, including developed pressure, +dP/dt, and -dP/dt returned to ~50% of baseline values at 45 min of reperfusion in both vehicle-treated WT and A<sub>3</sub>KO hearts, indicating that 20 min of global ischemia followed by reperfusion results in significant contractile dysfunction in our isolated mouse heart model system and that ischemic tolerance is similar between WT and A<sub>3</sub>KO mice. Treatment with 100 nM Cl-IB-MECA significantly improved reperfusion contractile function from ~50 to ~75% of baseline values in WT mouse hearts; this beneficial effect of Cl-IB-MECA was not apparent in hearts from A<sub>3</sub>KO mice (Fig. 7). In both WT and A<sub>3</sub>KO hearts, treatment with Cl-IB-MECA increased reperfusion coronary



**Fig. 7.** Recovery of left ventricular (LV) developed pressure (A), LV +dP/dt (B), LV -dP/dt (C), and coronary flow (D) expressed as a percentage of baseline of isolated Langendorff-perfused mouse hearts obtained from WT C57BL/6 or A<sub>3</sub>KO mice after 20 min of no-flow global ischemia and 45 min of reperfusion. Baseline parameters following equilibration for WT C57BL/6 and A<sub>3</sub>KO hearts, respectively, were LV dP = 140  $\pm$  5 and 149  $\pm$  6 mm Hg; LV +dP/dt = 6.7  $\pm$  0.8 and 6.3  $\pm$  0.8 mm Hg/ms; LV -dP/dt = 4.3  $\pm$  0.7 and 4.4  $\pm$  0.9 mm Hg/ms; and coronary flow = 23.1  $\pm$  1.5 and 22.3  $\pm$  2.0 ml/min/g. All data are presented as the mean  $\pm$  S.E.M. n = 8–12 hearts/group. \*, p < 0.05 versus the vehicle-treated group.

flow to preischemic levels. Collectively, these results suggest that cardioprotection provided by Cl-IB-MECA in isolated mouse hearts is the result of A<sub>3</sub>AR activation.


**Cardioprotection Provided by Cl-IB-MECA Is Not Mediated by the Release of Stored Mediators from Mast Cells.** Additional experiments were conducted to test the possibility that Cl-IB-MECA induces cardioprotection, because of its ability to stimulate mast cells, to degranulate. Theoretically, Cl-IB-MECA could provide cardioprotection by depleting mast cells of noxious mediators that contribute to injury during ischemia or induce adaptive changes that provide protection similar to that of myocardial preconditioning (Linden, 1994). To address this possibility, we conducted a series of experiments testing the cardioprotective effectiveness of Cl-IB-MECA in mice that had been treated chronically with compound 48/80 to deplete mast cells of stored mediators. Compound 48/80 was administered to WT mice intraperitoneally twice daily for 5 days beginning with a dose of 1 mg/kg on the first day, followed by an increment of 1 mg/kg per day, to a maximum of 5 mg/kg on the final day (Riley and West, 1955). On the following day (day 6), the effectiveness of Cl-IB-MECA was examined in both the in vivo mouse model of infarction and the isolated mouse heart model of global ischemia and reperfusion. As shown in Figs. 8 and 9, depletion of mast cell contents with compound 48/80 did not attenuate protection provided by Cl-IB-MECA (100  $\mu$ g/kg) in intact mice or in isolated mouse hearts. In the in vivo model, treatment with Cl-IB-MECA produced a 30% reduction in infarct size in compound 48/80-treated mice (Fig. 8). In the isolated mouse heart model, Cl-IB-MECA increased postischemic recovery of left ventricular developed pressure from 54 to 71% of preischemic baseline values after treatment with compound 48/80 (Fig. 9). Effective depletion of mast cell mediators was confirmed by the observation that

Cl-IB-MECA did not increase plasma histamine concentrations or reduce blood pressure in compound 48/80-treated mice (Fig. 8).

## Discussion

We observed that pretreatment with the A<sub>3</sub>AR-selective agonist Cl-IB-MECA produced a dose-dependent reduction in infarct size in mice subjected to 30 min of coronary artery occlusion and 24 h of reperfusion. This protective effect of Cl-IB-MECA was apparent at a dose of 30  $\mu$ g/kg and produced a  $\sim$ 35% reduction in infarct size at a dose of 100  $\mu$ g/kg. We subsequently observed that cardioprotection provided by Cl-IB-MECA was blocked completely in mice pretreated with the rodent A<sub>3</sub>AR antagonist MRS 1523 (Li et al., 1998), suggesting involvement of the A<sub>3</sub>AR. To further explore the cardioprotective profile of Cl-IB-MECA, we examined its effectiveness in an isolated mouse heart model of 20 min of normothermic global ischemia and 45 min of reperfusion. In hearts from WT mice, treatment with 100 nM Cl-IB-MECA significantly improved several indices of left ventricular postischemic contractile function including developed pressure; this protective effect of Cl-IB-MECA was lost completely using hearts obtained from A<sub>3</sub>KO mice. Collectively, these data are the first to provide definitive evidence that an A<sub>3</sub>AR agonist provides protection against myocardial ischemia/reperfusion injury by activating the A<sub>3</sub>AR. These data support the concept that the A<sub>3</sub>AR plays a protective role in the ischemic myocardium.

Several previous studies have demonstrated that A<sub>3</sub>AR agonists are effective at protecting against myocardial ischemia/reperfusion injury (Auchampach et al., 1997b, 2003; Tracey et al., 1997, 1998, 2003; Jordan et al., 1999; Thourani et al., 1999a,b; Kodani et al., 2001; Takano et al., 2001). For




**Fig. 8.** A, effect of Cl-IB-MECA on myocardial infarct size expressed as a percentage of the risk region induced by 30 min of LAD coronary occlusion and 24 h of reperfusion in WT C57BL/6 mice treated chronically with compound 48/80 to deplete mast cells of stored mediators. Vehicle or Cl-IB-MECA (100 µg/kg i.v.) was administered 10 min before the coronary occlusion. Risk region size was 39 ± 1 and 39 ± 2% of the left ventricle for the vehicle- and Cl-IB-MECA-treated groups, respectively. B and C, effect of Cl-IB-MECA (100 µg/kg) on mean arterial blood pressure and plasma histamine levels in compound 48/80-treated mice. Baseline mean arterial blood pressure was 68 ± 3 mm Hg. All data are presented as the mean ± S.E.M.  $n = 8$  (A) or 6–8 (B and C) mice/group. \*,  $p < 0.05$  versus the vehicle-treated group.

example, we have reported that pretreatment with IB-MECA effectively reduced infarct size, attenuated myocardial stunning, and induced the late phase of ischemic preconditioning in a clinically relevant conscious rabbit model or a barbital-anesthetized dog model of regional ischemia/reperfusion injury (Auchampach et al., 1997b, 2003; Kodani et al., 2001; Takano et al., 2001). In the dog model, IB-MECA was shown

to reduce infarct size even when administered immediately before release of the occlusion, demonstrating that it was also effective at reducing reperfusion-mediated injury (Auchampach et al., 2003). Similar results have been obtained by others using more selective A<sub>3</sub>AR agonists in in vivo models as well as in vitro models of ischemia/reperfusion injury, including isolated rodent hearts (Tracey et al., 1997, 1998, 2003; Thourani et al., 1999a,b). However, even though the A<sub>1</sub>AR was excluded using highly selective antagonists (Tracey et al., 1997, 1998, 2003; Kodani et al., 2001), involvement of the A<sub>3</sub>AR could not be proven definitively in these studies due to the lack of selective antagonists. Using the most recently developed rodent A<sub>3</sub>AR antagonist MRS 1523 and A<sub>3</sub>KO mice, we have now confirmed that protection against ischemia/reperfusion injury provided by Cl-IB-MECA in the mouse is mediated predominantly via A<sub>3</sub>AR.

We conducted additional experiments using ZM 241385 to address the possibility that Cl-IB-MECA may be producing cardiac protection by a mechanism involving the A<sub>2A</sub>AR. We chose to address this issue, because administration of low doses of A<sub>2A</sub>AR agonists has been reported to reduce myocardial reperfusion injury by suppressing inflammatory responses (Glover et al., 2004; Yang et al., 2005) and because it has been suggested that N<sup>6</sup>-benzyladenosine-5'-N-methylcarboxamide A<sub>3</sub>AR agonists may have higher affinity for the A<sub>2A</sub>AR than originally appreciated (Murphree et al., 2002). Moreover, Yang et al. (2004) have presented the hypothesis, in abstract form, that A<sub>3</sub>AR agonists may act by increasing adenosine production, which subsequently provides tissue protection via the A<sub>2A</sub>AR by an anti-inflammatory mechanism. ZM 241385 is a highly potent A<sub>2A</sub>AR antagonist with low affinity for the mouse A<sub>3</sub>AR (Palmer et al., 1995; Kreckler et al., 2006). Surprisingly, we found that pretreating mice with 2 mg/kg ZM 241385 (Ohta and Sitkovsky, 2001) was also capable of blocking the infarct size-reducing effect of Cl-IB-MECA in our in vivo mouse model. However, at this dose, ZM 241385 also inhibited the effect of Cl-IB-MECA to increase plasma histamine levels, a response in rodents known to be due to A<sub>3</sub>AR-mediated degranulation of mast cells (Linden, 1994; Van Schaik et al., 1996), confirming that a lower dose of ZM 241385 was required to prevent blockade of A<sub>3</sub>ARs. Using a dose of 0.5 mg/kg ZM 241385 that only partially inhibited Cl-IB-MECA-induced histamine release (but effectively blocked the hypotensive actions of the A<sub>2A</sub>AR agonist CGS 21680; Supplemental Fig. 1), we found that Cl-IB-MECA continued to produce a significant reduction in infarct size. These results exclude the involvement of the A<sub>2A</sub>AR in mediating the cardioprotective effects of Cl-IB-MECA and provide further evidence in support of an A<sub>3</sub>AR-mediated mechanism. Perhaps more importantly, these results demonstrate that ZM 241385 exhibits relatively high in vivo potency in rodents such that doses lower than 2 mg/kg should be used to selectively block A<sub>2A</sub>ARs.

We have previously reported in conscious rabbits and dogs that IB-MECA effectively reduced ischemia/reperfusion injury at doses that produced no changes in systemic hemodynamic parameters (Auchampach et al., 1997b, 2003; Kodani et al., 2001; Takano et al., 2001). This was an important observation, because the clinical use of adenosine as well as subtype-selective agonists for A<sub>1</sub> and A<sub>2A</sub>ARs for treating ischemia/reperfusion injury carry the risk of hemodynamic side-effects, including hypotension, bradycardia, and atrio-



**Fig. 9.** Recovery of left ventricular (LV) developed pressure (A), LV +dP/dt (B), LV -dP/dt (C), and coronary flow (D) expressed as a percentage of baseline of isolated Langendorff-perfused mouse hearts obtained from WT C57BL/6 mice treated with compound 48/80 to deplete mast cell of stored mediators. Baseline parameters following equilibration were LV dP = 168 ± 7 mm Hg; LV +dP/dt = 7.7 ± 0.4 mm Hg/ms; LV -dP/dt = 3.6 ± 0.2 mm Hg/ms; and coronary flow = 28.9 ± 3.3 ml/min/g. All data are presented as the mean ± S.E.M.  $n = 8$  hearts/group. \*,  $p < 0.05$  versus the vehicle-treated group.

ventricular blockade, due to activation of A<sub>1</sub>ARs expressed in conducting tissue and A<sub>2A</sub>ARs in vascular smooth muscle cells. However, in rodent species (but not in other species), it is well known that activation of A<sub>3</sub>ARs reduces blood pressure due to the actions of vasoactive mediators released from mast cells (Linden, 1994; Hannon et al., 1995; Fozard et al., 1996). Indeed, in the present investigation, we observed that administration of 100  $\mu$ g/kg Cl-IB-MECA produced an immediate (within 1–2 min) 10 to 15% reduction in blood pressure, which correlated with increased plasma histamine concentrations, but was absent in compound 48/80-treated mice. Interestingly, however, we observed that the kinetic profile of Cl-IB-MECA to reduce blood pressure differed between the two strains of mice that we used in our studies, in that the response was transient in FVB/N mice (~15–20 min) but more sustained (at least 45 min) in C57BL/6 mice. Because Cl-IB-MECA continued to produce a delayed reduction in blood pressure in WT or A<sub>3</sub>KO mice pretreated with MRS 1523 or a nonselective dose of ZM 241385 (2 mg/kg), we conclude that Cl-IB-MECA reduced blood pressure in C57BL/6 mice by a mechanism that may not be receptor-dependent.

We did not address the specific mechanism by which Cl-IB-MECA exerted cardioprotection, although we did determine that it is independent of mast cell degranulation. We have previously shown that the infarct size-reducing effects of IB-MECA are blocked by glibenclamide (Auchampach et al., 1997b, 2003), implicating involvement of ATP-sensitive potassium channels. We have also determined in preliminary studies using our in vivo mouse model of infarction and A<sub>3</sub>KO bone marrow chimeric mice that Cl-IB-MECA seems to re-

duce injury when administered at the time of reperfusion by an effect on bone marrow-derived cells, suggesting an anti-inflammatory mechanism (Ge et al., 2004). Therefore, we predict that A<sub>3</sub>AR agonists function by different mechanisms to protect against ischemia/reperfusion injury depending on whether they are administered before ischemia or during reperfusion. Although it is unlikely to be a significant factor, since we observed protection in the isolated heart model and in compound 48/80-treated mice, it is important to note that the hypotensive action produced by Cl-IB-MECA in the present investigation may have contributed to the reduction in infarct size in the in vivo studies.

One aspect of our ischemia/reperfusion studies using A<sub>3</sub>KO mice requires particular attention. In an earlier study, we reported that infarct size was smaller in A<sub>3</sub>KO mice subjected to 30 min of coronary artery occlusion and 24 h of reperfusion, suggesting that deletion of the A<sub>3</sub>AR gene induces an unexpected cardioprotective phenotype (Guo et al., 2001). However, this earlier study was conducted with A<sub>3</sub>KO mice on a mixed background of three different genetic strains (C57BL/6, sv129, and D2). Furthermore, controls used in this previous study were not littermates but were from a separate line of mice interbred among the three different genetic strains. This line of control mice was considered to be the best available at that time. Based on the results of the present investigation using congenic (C57BL/6) A<sub>3</sub>KO mice, we conclude that genetic deletion of the A<sub>3</sub>AR gene itself has no effect on ischemic tolerance and that the results of our earlier study are probably explained by differences in the genetic background of the mice rather than specific deletion of the

$A_3$ AR gene (Guo et al., 2001). Harrison et al. (2002) reached a similar conclusion regarding  $A_3$ KO mice.

In summary, we have demonstrated that pretreatment with the  $A_3$ AR agonist Cl-IB-MECA protects against ischemia/reperfusion injury in mice and that this occurs through an  $A_3$ AR-mediated mechanism that is independent of mast cell degranulation. These results further support the contention that therapeutic strategies targeting the  $A_3$ AR could be a novel and useful approach for protection of the ischemic myocardium.

#### Acknowledgments

We acknowledge the Cardiovascular Center HPLC Core Facility at the Medical College of Wisconsin for purification of radioligands.

#### References

Auchampach JA, Ge ZD, Wan TC, Moore J, and Gross GJ (2003) The  $A_3$  adenosine receptor agonist IB-MECA reduces myocardial ischemia/reperfusion injury in dogs. *Am J Physiol* **285**:H607–H613.

Auchampach JA, Jin X, Wan TC, Caughey GH, and Linden J (1997a) Canine mast cell adenosine receptors: cloning and expression of the  $A_3$  receptor and evidence that degranulation is mediated by the  $A_{2B}$  receptor. *Mol Pharmacol* **52**:846–860.

Auchampach JA, Rizvi A, Qiu Y, Tang XL, Maldonado C, Teschner S, and Bolli R (1997b) Selective activation of  $A_3$  adenosine receptors with  $N^6$ -(3-iodobenzyl)adenosine-5'-N-methyluronamide protects against myocardial stunning and infarction without hemodynamic changes in conscious rabbits. *Circ Res* **80**:800–809.

Black RG Jr, Guo Y, Ge ZD, Murphree SS, Prabhu SD, Jones WK, Bolli R, and Auchampach JA (2002) Gene dosage-dependent effects of cardiac-specific overexpression of the  $A_3$  adenosine receptor. *Circ Res* **91**:165–172.

Cross HR, Murphy E, Black RG, Auchampach J, and Steenbergen C (2002) Overexpression of  $A_3$  adenosine receptors decreases heart rate, preserves energetics, and protects ischemic hearts. *Am J Physiol* **283**:H1562–H1568.

Fozard JR, Pfannkuche HJ, and Schuurman HJ (1996) Mast cell degranulation following adenosine  $A_3$  receptor activation in rats. *Eur J Pharmacol* **298**:293–297.

Ge ZD, Wan TC, and Auchampach JA (2004) The  $A_3$  adenosine receptor agonist Cl-IB-MECA reduces myocardial infarct size in mice when administered during reperfusion: mechanistic studies with  $A_3$ AR gene 'knock-out' and bone marrow chimeric mice. *Circulation* **110**:III-29.

Glover DK, Riou LM, Ruiz M, Sullivan GW, Linden J, Rieger JM, Macdonald TL, Watson DD, and Beller GA (2005) Reduction of infarct size and postischemic inflammation from ATL-146e, a highly selective adenosine  $A_{2A}$  receptor agonist, in reperfused canine myocardium. *Am J Physiol* **288**:H1851–H1858.

Guo Y, Bolli R, Bao W, Wu WJ, Black RG Jr, Murphree SS, Salvatore CA, Jacobson MA, and Auchampach JA (2001) Targeted deletion of the  $A_3$  adenosine receptor confers resistance to myocardial ischemic injury and does not prevent early preconditioning. *J Mol Cell Cardiol* **33**:825–830.

Hannon JP, Pfannkuche HJ, and Fozard JR (1995) A role for mast cells in adenosine  $A_3$  receptor-mediated hypotension in the rat. *Br J Pharmacol* **115**:945–952.

Harrison GJ, Cerniway RJ, Peart J, Berr SS, Ashton K, Regan S, Paul Matherne G, and Headrick JP (2002) Effects of  $A_3$  adenosine receptor activation and gene knock-out in ischemic-reperfused mouse heart. *Cardiovasc Res* **53**:147–155.

Jordan JE, Thourani VH, Auchampach JA, Robinson JA, Wang NP, and Vinten-Johansen J (1999)  $A_3$  adenosine receptor activation attenuates neutrophil function and neutrophil-mediated reperfusion injury. *Am J Physiol* **277**:H1895–H1905.

Kodani E, Bolli R, Tang XL, and Auchampach JA (2001) Protection of IB-MECA against myocardial stunning in conscious rabbits is not mediated by the  $A_1$  adenosine receptor. *Basic Res Cardiol* **96**:487–496.

Kreckler LM, Wan TC, Ge ZD, and Auchampach JA (2006) Adenosine inhibits TNF- $\alpha$  release from mouse peritoneal macrophages via  $A_{2A}$  and  $A_{2B}$ , but not  $A_3$  adenosine receptors. *J Pharmacol Exp Ther* **317**:172–180.

Lappas CM, Sullivan GW, and Linden J (2005) Adenosine  $A_{2A}$  agonists in development for the treatment of inflammation. *Expert Opin Investig Drugs* **14**:797–806.

Li AH, Moro S, Melman N, Ji XD, and Jacobson KA (1998) Structure-activity relationships and molecular modeling of 3,5-diacyl-2,4-dialkylpyridine derivatives as selective  $A_3$  adenosine receptor antagonists. *J Med Chem* **41**:3186–3201.

Linden J (1994) Cloned adenosine  $A_3$  receptors: pharmacological properties, species differences and receptor functions. *Trends Pharmacol Sci* **15**:298–306.

Murphree LJ, Marshall MA, Rieger JM, MacDonald TL, and Linden J (2002) Human  $A_{2A}$  adenosine receptors: high-affinity agonist binding to receptor-G protein complexes containing  $G_b$ . *Mol Pharmacol* **61**:455–462.

Ohta A and Sitkovsky M (2001) Role of G-protein-coupled adenosine receptors in downregulation of inflammation and protection from tissue damage. *Nature (Lond)* **414**:916–920.

Olah ME, Gallo-Rodriguez C, Jacobson KA, and Stiles GL (1994)  $^{125}$ I-4-Aminobenzyl-5'-N-methylcarboxamidoadenosine, a high affinity radioligand for the rat  $A_3$  adenosine receptor. *Mol Pharmacol* **45**:978–982.

Palmer TM, Poucher SM, Jacobson KA, and Stiles GL (1995)  $^{125}$ I-4-(2-[7-Amino-2-[2-furyl][1,2,4]triazolo[2-a][1,3,5]triazin-5-yl-amino]ethyl)phenol, a high affinity antagonist radioligand selective for the  $A_{2A}$  adenosine receptor. *Mol Pharmacol* **48**:970–974.

Peart J, Flood A, Linden J, Matherne GP, and Headrick JP (2002) Adenosine-mediated cardioprotection in ischemic-reperfused mouse heart. *J Cardiovasc Pharmacol* **39**:117–129.

Riley JF and West GB (1955) Tissue mast cell studies with a histamine-liberator with low toxicity (compound 48/80). *J Pathol Bacteriol* **69**:269–282.

Salvatore CA, Tilley SL, Latour AM, Fletcher DS, Koller BH, and Jacobson MA (2000) Disruption of the  $A_3$  adenosine receptor gene in mice and its effect on stimulated inflammatory cells. *J Biol Chem* **275**:4429–4434.

Takano H, Bolli R, Black RG Jr, Kodani E, Tang XL, Yang Z, Bhattacharya S, and Auchampach JA (2001)  $A_1$  or  $A_3$  adenosine receptors induce late preconditioning against infarction in conscious rabbits by different mechanisms. *Circ Res* **88**:520–528.

Thourani VH, Nakamura M, Ronson RS, Jordan JE, Zhao ZQ, Levy JH, Szlam F, Guyton RA, and Vinten-Johansen J (1999a) Adenosine  $A_3$ -receptor stimulation attenuates postischemic dysfunction through  $K_{ATP}$  channels. *Am J Physiol* **277**:H228–H235.

Thourani VH, Ronson RS, Jordan JE, Guyton RA, and Vinten-Johansen J (1999b) Adenosine  $A_3$  pretreatment before cardioplegic arrest attenuates postischemic cardiac dysfunction. *Ann Thorac Surg* **67**:1732–1737.

Tracey WR, Magee W, Masamune H, Kennedy SP, Knight DR, Buchholz RA, and Hill RJ (1997) Selective adenosine  $A_3$  receptor stimulation reduces ischemic myocardial injury in the rabbit heart. *Cardiovasc Res* **33**:410–415.

Tracey WR, Magee W, Masamune H, Oleynek JJ, and Hill RJ (1998) Selective activation of adenosine  $A_3$  receptors with  $N^6$ -(3-chlorobenzyl)-5'-N-methylcarboxamidoadenosine (CB-MECA) provides cardioprotection via  $K_{ATP}$  channel activation. *Cardiovasc Res* **40**:138–145.

Tracey WR, Magee WP, Oleynek JJ, Hill RJ, Smith AH, Flynn DM, and Knight DR (2003) Novel  $N^6$ -substituted adenosine 5'-N-methyluronamides with high selectivity for human adenosine  $A_3$  receptors reduce ischemic myocardial injury. *Am J Physiol* **285**:H2780–H2787.

Van Schaik EA, Jacobson KA, Kim HO, IJzerman AP, and Danhof M (1996) Hemodynamic effects and histamine release elicited by the selective adenosine  $A_3$  receptor agonist 2-Cl-IB-MECA in conscious rats. *Eur J Pharmacol* **308**:311–314.

Yang Z, Day YJ, Toufektsian MC, Ramos SI, Marshall M, Wang XQ, French BA, and Linden J (2005) Infarct-sparing effect of  $A_{2A}$ -adenosine receptor activation is due primarily to its action on lymphocytes. *Circulation* **111**:2190–2197.

Yang Z, Marshal M, Xu Y, French BA, and Linden J (2004) Opposing effects of the adenosine receptor agonist IB-MECA on myocardial infarct size in mice are mediated by  $A_3$  proinflammatory receptors and anti-inflammatory  $A_{2A}$  receptors. *Circulation* **110**:III-300.

**Address correspondence to:** Dr. John A. Auchampach, Department of Pharmacology and Toxicology, 8701 Watertown Plank Road, Medical College of Wisconsin, Milwaukee, WI 53226. E-mail: jauchamp@mcw.edu