

Utah State University
DigitalCommons@USU

All Graduate Theses and Dissertations

Graduate Studies

5-2010

Comparative Performance of Fluorometry and High Performance Liquid Chromatography in the Detection of Aflatoxin M1 in Two Commercial Cheeses

Gustavo Peña
Utah State University

Follow this and additional works at: <https://digitalcommons.usu.edu/etd>

 Part of the [Agronomy and Crop Sciences Commons](#), [Animal Sciences Commons](#), and the [Food Science Commons](#)

Recommended Citation

Peña, Gustavo, "Comparative Performance of Fluorometry and High Performance Liquid Chromatography in the Detection of Aflatoxin M1 in Two Commercial Cheeses" (2010). *All Graduate Theses and Dissertations*. 735.

<https://digitalcommons.usu.edu/etd/735>

This Thesis is brought to you for free and open access by the Graduate Studies at DigitalCommons@USU. It has been accepted for inclusion in All Graduate Theses and Dissertations by an authorized administrator of DigitalCommons@USU. For more information, please contact digitalcommons@usu.edu.



COMPARATIVE PERFORMANCE OF FLUOROMETRY AND HIGH
PERFORMANCE LIQUID CHROMATOGRAPHY IN
THE DETECTION OF AFLATOXIN M₁
IN TWO COMMERCIAL CHEESES

by

Gustavo Peña

A thesis submitted in partial fulfillment
of the requirements for the degree

of

MASTER OF SCIENCE

in

Dairy Science

Approved:

Allen J. Young
Major Professor

Jeffery O. Hall
Committee Member

Jong-Su Eun
Committee Member

Kenneth L. White
Department Head

Byron R. Burnham
Dean of Graduate Studies

UTAH STATE UNIVERSITY
Logan, Utah

2010

Copyright © Gustavo Pena

All Rights Reserved

ABSTRACT

Comparative Performance of Fluorometry and High Performance Liquid Chromatography in the
Detection of Aflatoxin
M₁ in Two Commercial Cheeses

by

Gustavo Peña, Master of Science

Utah State University, 2009

Major Professor: Dr. Allen Young
Department: Animal, Dairy and Veterinary Sciences

Aflatoxin M₁ (**AFM₁**) is frequently found in milk and dairy products. It is a metabolite formed in cows from aflatoxin B₁ (**AFB₁**), contained in animal feeds. In cheese production AFM₁ distributes between curds and whey. In this study, cows were fed 64 µg/AFB₁/d for the high treatment, and 5 µg/AFB₁/d for the low treatment, to obtain milk contaminated with AFM₁ over the 0.5 µg/L and under 0.05 µg/L restrictions, respectively. Cheese was manufactured with milk contaminated with AFM₁ at 0.8 and 0.03 µg/kg by the higher and lower treatment, respectively. Two commercial cheeses were elaborated: a hard-aged cheese (cheddar cheese) and soft high moisture cheese (fresco cheese) to evaluate whether the cheese type had any impact on AFM₁ analysis. AFM₁ was extracted from cheese using immunoaffinity columns. Analyses were carried out by using high pressure liquid chromatography (**HPLC**) as the reference method and fluorometry as a method of validation. Analysis was by 2-way fixed factor analyses. AFM₁ was detected in all samples by both methods of analysis. There were no detectable statistical

differences between cheese types ($P>0.05$). AFM₁ content was significantly different between the high and low concentration of AFB₁ used to make the cheese type ($P<0.01$). Our regression model shows a linear relationship between fluorometry and HPLC methods; $R^2=0.9141$ from cheddar cheese and $R^2=0.9141$ from fresco cheese. There were no statistical differences between methods of analysis ($P>0.05$). Carryover of AFM₁ in cheese detected by fluorometry in cheddar cheese was 163% and 80% for high and low treatments, respectively, and in fresco cheese was 119 and 133 for high and low treatments, respectively. These carryovers are below that reported in the literature. Results suggest that fluorometry is a simple and reliable AFM₁ detection method for screening samples of complex matrices such as cheese.

(58 pages)

ACKNOWLEDGMENTS

I will always be grateful for the chance given to pursue a graduate education. I wish to express my sincere gratitude and appreciation to the people who have both directly and indirectly contributed to this work. The following are those to whom I am particularly indebted. I would like to thank:

My major professor, Dr. Allen Young, for his patience, and my committee members, Dr. Jeffery O. Hall and Dr. Jong-Su Eun, for their assistance during the performance of my dissertation work. Special recognition goes to Dr. Duarte Diaz for his actions and assistance in my graduate research project. It was a great experience working and traveling with him. Dr. Antonio Gallo and Pam Hole are thanked for their useful advice, assistance, the use of equipment and many good times during my graduate work.

Special thanks go to the USU-Caine Dairy Farm and Nutrition, Dietetics, Food and Science Department and staff for their support, assistance, advice, and patience in my attempts to milk cows and make cheese. I would like at this time to dedicate this thesis to my family, especially my parents, Florentino and Antonia Peña. I love you both very much. My wife, Elvia, and my sons Eduardo and Zuzet are specially recognized for their support, patience, tolerance, and understanding. I would like to thank my brothers and sisters for their encouragement and support through my graduate work. To the rest of my family and friends-thanks for your encouragement.

G. Peña

CONTENTS

	Page
ABSTRACT	iii
ACKNOWLEDGMENTS	v
LIST OF TABLES	viii
LIST OF FIGURES	ix
INTRODUCTION	1
LITERATURE REVIEW	3
Mold Growth and Mycotoxin Formation.....	3
Aflotoxins	4
Worldwide Occurrence of Aflatoxin M ₁	6
Biological Action of Aflatoxin	6
Effects of Aflatoxin in Ruminants	9
Effects of Aflatoxin in Humans	11
Aflotoxin Analysis	13
Sample Selection and Preparation	14
Extraction	14
Clean-up	15
Detection	15
Fluorometry Analysis.....	16
High Pressure Liquid Chromatography Analyses.....	18
Aflatoxin in Milk	21
Aflatoxin Determination in Cheese	21
Control and Prevention of Aflatoxin M ₁	22
Developing Methods for Aflatoxin M ₁ in Cheese	24
MATERIALS AND METHODS.....	26
Contaminated Corn with Aflatoxin B ₁	26
Animals	26
Milk Collection and Processing	27
Cheese Production	28
Making of Cheddar Cheese.....	29
Making of Fresco Cheese.....	29

Detection and Quantification of Aflatoxin M ₁ from Cheese	30
Chemical, Standards and Laboratory Supply.....	30
Extraction of Aflatoxin M ₁ from Cheese.....	30
Aflatoxin M ₁	31
Florometry Determination	31
High Pressure Liquid Chromatography	32
Stability of the Methods.....	33
Statistical Analysis.....	33
RESULTS AND DISCUSSION	34
Carryover of Aflatoxin M ₁ Feed Trail	34
Detection of Aflatoxin M ₁ in Cheese.....	35
Stability of the Analytical Method.....	37
Fluorometry Cross-Validation	38
..... 1Agricultural Education for Students with Disabilities	
..... 16 Summary	
..... 19	
Estimation of the Precision	39
Demonstration of Accuracy	39
Comparison of the Results Obtained by the Two Methods	39
CONCLUSION.....	40
..... 1Agricultural Education for Students with Disabilities	
..... 16Summary	
..... 19	
REFERENCES	43

LIST OF TABLES

Table		Page
1	Comparison of Physiobiochemical Parameters for Fungi Growth and Aflatoxin Production of <i>Aspergillus flavus</i> and <i>A. parasiticus</i>	4
2	Data of the Occurrence of Aflatoxin M ₁ in Cheese on Several Countries	7
3	A Review of Methods Available for Mycotoxin Analyses	16
4	Ingredients and Chemical Composition of the Diet	27
5	Concentration of Aflatoxin M ₁ in Milk Prior to Cheese Making. Evaluation was Made April 16, 2007 by the USU Lab and February 12, 2009 by the External Lab.....	34
6	Means of Aflatoxin M ₁ Means for High Pressure Liquid Chromatography and Fluorometry Method from Different Cheese Type and Aflatoxin M ₁ Fed Concentrations.....	35
7	Mean Carryover of Aflatoxin M ₁ in Cheese as Compared to Milk Content.....	36
8	Stability of the Analytical Method for Aflatoxin M ₁ Detected by Fluorometry and HPLC.....	39
9	Comparison of Variables from Fluorometry against High Pressure Liquid Chromatography.....	40

LIST OF FIGURES

Figure	Page
1 Relationship of Aflatoxin M ₁ High and Low Concentration between HPLC and Fluorometry Method from Cheddar Cheese	38
2 Relationship of Aflatoxin M ₁ High and Low Concentration between HPLC and Fluorometry Method from Fresco Cheese.....	38

INTRODUCTION

The ubiquitous nature of fungi (Coulombe, 1993), creates a worldwide problem by contaminating food and feeds (Kamkar, 2005). Such a problem is exacerbated by change in climatic conditions, agricultural manufacturing, transportation and storage practices (CAST, 2003). Some varieties of fungi are capable of produce secondary metabolites called mycotoxins; today more than 100 molds are known to be responsible for producing approximately 400 mycotoxins with toxigenic potential (Kabak et al., 2006). In domestic animals, ingestion of this toxic compounds compromise economic and productive parameters (Binder et al., 2007). In addition, many mycotoxins represent a threat to human health (Coulombe, 1993).

Aflatoxins (AF) are secondary metabolites produced by five species of the fungi *Aspergillus*: *A. flavus*, *A. Parasiticus*, *A. nomius*, *A. tamari* and *A. pseudotamarii*. Only *A. flavus* and *A. parasiticus* produce high enough concentration to be economically important. Aflatoxins are produced by the fungi in response to changes in environmental conditions, contaminating the substrates on what they grow either pre-harvest or during post-harvest storage (Flannigan, 1991; Kozakiewicz and Smith, 1994).

Aspergillus mainly affects corn, peanuts, cotton, tree nuts and their by-products (CAST, 2003). In the US, the Food and Drug Administration regulates AFB₁ in feeds intended for dairy cattle consumption at 20 µg/kg and limits AFM₁ in milk for human consumption at 0.5 µg/L (Coulombe, 1993).

Aflatoxins are the most studied group of mycotoxins because of their natural carcinogenic properties (Hussain and Anwar, 2007). Aflatoxin B₁ (AFB1) ingested from

feed is metabolized to aflatoxin M₁ (AFM₁) and excreted in milk and urine. According to the International Agency of Research on Cancer (IRCA), AFB₁ and AFM₁ have been classified as human carcinogens (Hussain and Anwar, 2007).

Quantitative carryover of AFB₁ to milk is around 3% of the concentration consumed and it is excreted into milk primarily as AFM₁ (Diaz et al., 2004). When contaminated milk is used to manufacture dairy products the toxins is transferred to the final product. The AFM₁ is unaffected by pasteurization or processing (Galvano et al., 2005). Studies have demonstrated that AFM₁ binds mainly with casein increasing the concentration in the curd during cheese manufacturing (Galvano et al., 2005).

Sensitive and reliable methods have been developed for detection in order to control the AFM₁ concentration in cheese. High Performance Liquid Chromatography (HPLC) analysis is one of the best choices when sensitivity, reliability and precision are required (Scudamore, 2005). More rapid methods like flourometry are being developed to determine AFM₁ and AFB₁ in foods, but there are not publications to investigate the use of fluorometry in complex foods like cheese.

In this study, the objective was to evaluate the efficiency of a quick analytical fluorometry procedure against the HPLC method at two different concentrations of AFM₁ residues in two kinds of cheeses (Fresco and Cheddar cheese), while studying the carryover of AFM₁ to cheese and the recovery AFM₁ using immunoaffinity columns as a clean-up step.

LITERATURE REVIEW

Mold Growth and Mycotoxin Formation

With a family of over 200 members, *Aspergillus* molds were first described in 1729 by an Italian priest, Micheli, who took the name from the shape of an aspergillum (holy water sprinkler) (Samson, 1992). Today those fungi are recognized as the most common genera in the world. *Aspergillus* was the first mold organism cultivated in artificial media for their biochemical properties. In Asian countries, *Aspergillus spp* have been used to ferment foods (Gervais and Bensoussan, 1994). They are known for the many contributions to biochemistry, molecular biology, serology and recently their potential as mycotoxin producers (Samson, 1992).

The genera *Aspergillus* is a soil born fungus important for its ability to break down vegetative material (CAST, 2003). However, a small group of *Aspergillus* species are mycotoxin producers. Species of the complex genera, *Aspergillus* has been adapted to a wide range of temperatures and habitats. They are most abundant in subtropical and warm climates (Krska, 1999; CAST, 2003). Various species are involved in producing AF on grains in the field or under storage conditions; in most parts of the world, *Aspergillus flavus* and *A. parasiticus* are the most representative agents although other species such as *A. nomius*, and *Penicilium* may be implicated (Agag, 2004).

Filament molds like *Aspergillus* are affected by physiobiochemical parameters such as water, temperature, pH and gas composition. Fungi cannot germinate without water but growth rate is controlled by temperature; such parameters are different for optimum mycotoxin formation than for optimal fungus growth (Kozakiewicz and Smith,

1994). (See Table 1.) Water activity (a_w) below 0.70 may affect germination and mycelial growth and higher than 0.98 increases the lag time before germination from a few days (Kozakiewicz and Smith, 1994). Production of AFB1 ranges from 12 to 41 °C with most favorable temperature occurs between 24 to 28 °C (Kozakiewicz and Smith, 1994; Agag, 2004). Growth pH is difficult to control because fungi change the pH in the substrate when growing. Some growth inhibitions have been observed by *Aspergillus* after O₂ concentration was below 1% (Kozakiewicz and Smith, 1994). However there are complex interactions between those parameters (Kozakiewicz and Smith, 1994).

Table 1. Comparison of Physiobiochemical Parameters for Fungus Growth and Aflatoxin Production of *Aspergillus flavus* and *A. parasiticus*¹

	a_w ²	Moisture % ³	Temperature ⁴	pH	CO ₂ Gas	Stress
Growth	0.78	18 - 25	25 – 37 °C	4.0 – 6.5	79 %	Reduced
Toxin ⁵	0.82		12 – 41 °C			Increased

¹ Modified from (Kozakiewicz and Smith, 1994; Agag, 2004).

² a_w = water activity for optimal growth and mycotoxin production

³ Field = 22 – 25; Storage = 13 – 18; Decay 18

⁴ Optimum temperatures for toxin production range from 25 – 32 °C

⁵ Toxin production

Aflatoxins

The first described report of aflatoxicosis was in a British journal in 1960 where a strange disease, “Turkey “X” disease,” was affecting and killing turkeys and other animals at an early age (Sargent et al., 1961). The disease was not cured by antibiotic treatment, and no bacterial, viral or pesticide agent was isolated (Carnaghan and Sargeant, 1961). The toxic compound was named aflatoxin from letters in the name *Aspergillus flavus* and the word toxin: “(a- (*aspergillus*) + fla-(*flavus*) + toxin)”. The

fungus and the toxin were found in peanut meal coming from Brazil (Sargent et al., 1961).

When fungi grows in cereal or feedstuff used to feed dairy cattle, there is a chance that mycotoxins will end up in milk and their carryover represents a latent threat to the health of the human population (Coffey et al., 2009). The production of AFB₁ and aflatoxin B₂ (AFB₂) is attributable to *A. flavus*, while *A. parasiticus* produces AFB₁, AFB₂, aflatoxin G₁ (AFG₁) and aflatoxin G₂ (AFG₂) (Agag, 2004). The letters B and G were from the fluorescense color produced under ultraviolet light (B for blue and G for green). Further metabolisms of AFB₁ and AFB₂ in animals produce the hydroxylated metabolites AFM₁ and aflatoxin M₂ (AFM₂), which are found in urine and milk. Although 18 types of AF are known, the most toxic are AFB₁ and AFM₁. The following metabolites are listed in order of decreased toxicity: AFG₁, AFM₂, AFB₂, AFG₂ (Gimeno and Martins, 2006).

Aflatoxin occurs as natural contaminants in diverse substrates. They are mainly found in cereals like corn, wheat and rice, cereal byproducts, peanuts, pistachio and other nuts, cottonseed, oilseed meals, cassava, dry fruits, sausage products, spices, wine, coffee, legumes, fruits and their juices, milk and dairy products (Flannigan, 1991). While the four naturally occurring AF (B₁, B₂, G₁, G₂) are common in the same foods, AFB₁ represents the highest percentage (60 to 80%) of the total aflatoxin content (Weidenborner, 2001).

Despite the fact that the liver is the target organ for AF, tumors in the respiratory systems have been associated with AF intoxication. The hepatotoxicity, immunodeficiency, acute and chronic toxicity will be discussed in later sections.

Worldwide Occurrence of Aflatoxin M₁

The problem of food contamination with AFM₁ is of great concern and has received enormous attention. The International Cancer Research Association classified AFB₁ as a Group 1 carcinogen (known carcinogen to humans) while AFM₁ is classified as a Group 2B carcinogen (possibly carcinogenic to humans) (IACR, 1997). Because of this, many countries today have been implementing controls and inspection programs to monitor the presence of AF in feeds and foods, resulting in maximum tolerance law for such compounds (FAO, 2004).

Milk and dairy products are the main commodities to introduce AF, primarily AFM₁ into the human diet. Many European countries have imposed a limit of 0.05 µg/L of AFM₁ in milk. However, in the US and other Latin American countries the maximum concentration is 0.5 µg/L. By 2003 AFM₁ regulation in food was undertaken by 98 countries (FAO, 2004). Data from a recent publication on the world wide occurrences of AFM₁ in cheese are summarized in Table 2. As expected there are more reports from regions where climate is favorable to AF production. In the U.S., the only report of milk contaminated by AFM₁ was in 1978 in Arizona where 413, 636 kg of milk were discarded (Park, 1993).

Biological Action of Aflatoxin

The biosynthesis of AF is the best characterized of any mycotoxin (CAST, 2003). Aflatoxin represents a group of difuranocoumarin compounds classified into two groups: the difurocoumarocyclopentenone which include AFB₁, AFB₂, aflatoxin B_{2a} (AFB_{2a}), AFM₁, AFM₂, aflatoxin M_{2a} (AFM_{2a}) and aflatoxicol (AFL). The other group,

Table 2. Data of the Occurrence on Aflatoxin M₁ in Cheese on Several Countries

Country	Sample	Positive	Method ³	Range, ng/kg	Reference
Brazil	75	56 ¹ (75%) ²	HPLC	20 - 6920	Prado, 2000
Libya	20	15 ¹ (75 %) ²	HPLC	110 - 520	Elgerbi, 2004
Turkey	63	28 ¹ (44 %) ²	ELISA	7 – 202	Gurses, 2004
Turkey	400	327 ¹ (82 %) ²	ELISA	> 250	Sarlmehtmetoglu, 2004

¹Represents number of positive samples from all samples

²Represents the percentage of positive samples

³HPLC = high performance liquid chromatography; ELISA = Enzyme linked immunosorbent assay

difurocoumarolactone is represented by AFG₁, AFG₂, aflatoxin G_{2a} (AFG_{2a}), aflatoxin GM₁ (AFGM₁), aflatoxin GM₂ (AFGM₂), aflatoxin GM_{2a} (AFGM_{2a}), and aflatoxin B₃ (AFB₃) (Agag, 2004).

Following a rapid absorption from the gastrointestinal tract, AFB₁ tends to accumulate in the liver. Metabolism is mainly in the liver but other tissues are also involved (Agag, 2004). Several metabolites can be produced, some of which are highly reactive and can react with cellular macromolecules. The reactive metabolites attach to proteins and nucleic acids. Unbound metabolites are excreted in urine and milk, and conjugated aflatoxin is excreted in feces (Riley and Pestka, 2005).

Aflatoxins enter the cell and are metabolized via monooxygenases in the endoplasmatic reticulum to hydroxylated metabolites which are further metabolized to glucuronide and sulfate conjugates or oxidized to the reactive epoxide which undergoes spontaneous hydrolysis to AFB₁-8,9-dihydrodiol or bind to proteins, resulting in cytotoxicity. The epoxide can react with DNA or protein, or be detoxified by an inducible

GST to the GSH-conjugate. Both DNA and protein adducts have been proven to be useful as biomarkers in humans and laboratory animals to monitor AF exposure (Riley and Pestka, 2005).

Toxic effects of AF are related to their biotransformation pathway (Eaton and Gallagher, 1994). In the liver and other tissues, AFB₁ requires microsomal oxidation to the reactive AFB₁-8, 9-epoxide to produce its hepatotoxicity and carcinogenic effects. The oxidation also produces other polar metabolites like AFM₁. During the epoxidation of AFB₁, microsomal cytochrome P450 (CYP450) generates a reactive epoxide responsible for nucleic acid alkylation. Even though CYP450 is the main pathway to form epoxide, independent pathways have also been demonstrated (Eaton and Gallagher, 1994).

Monoxygenase hydroxylation of AFB₁ produces several metabolites (aflatoxin Q₁ (AFQ₁), aflatoxin P₁ (AFP₁) and AFM₁) with lower carcinogenic potential than parent toxin. AFQ₁ is created via 3a-hydroxylation. Aflatoxin P₁ is formed by o-demethylation and AFM₁ is formed by 9a-hydroxylation. AF can be reduced to AFL by reduction of the 1-keto-group through a cytosolic NADPH-dependent reductase. The formations of AFL may not be an excretion pathway because AFL rapidly converts back to AF by microsomal dehydrogenase. Another product of reduction is the AFB₁-8,9-aldehyde formed by hydrolysis of AFB₁-8,9-epoxide. This may be partially responsible for the acute toxic effects of AFB₁ (Eaton and Gallagher, 1994).

Detoxification of AF epoxide is mediated by glutation-S-transferase (GST) by conjugation with reduced glutathione (GSH) in the liver. Other conjugation pathways include glucuronidation and sulfations of hydroxylated metabolites are excreted in biliary

fluids. AFQ₁ and AFP₁ may conjugate with GSH to form detoxification products (Eaton and Gallagher, 1994).

Aflatoxin M₁ is a metabolite found in milk of animals consuming AF contaminated feed. According to Diaz et al. (2004), the AFM₁ hydroxilated metabolite of AFB₁ can be detectable during the first milking after animals consumed AF contaminated feed. It reached its' maximum concentration in three days and disappeared four days after the last feeding time of contaminated feed. AFM₁ has lower carcinogenic potency than AFB₁, with carcinogenic properties of AFM₁ being 30% less in trout than AFB₁ (Eaton and Gallagher, 1994). Nuryono et al. (2008) cited authors who found that AFM₁ is about 10 times less carcinogenic than AFB₁, but its genotoxicity is higher than the parent toxin. Even with lower toxicity, AFM₁ is hepatotoxic and carcinogenic (Coulombe, 1993).

Effects of Aflatoxin in Ruminants

Detrimental effect of AF in animals by damage to vital organs causes serious economic impact (Coulombe, 1993; Whitlow, 2005). Aflatoxin is known to cause liver damage and immmunosuppresion, as well as having carcinogenic, teratogenic and mutagenic effects. But, vulnerability is influenced by breed, species, age, dose, length of exposure and nutritional status of the animals (Richard, 2007). Calves that consume AFM1 from milk are more susceptible than older animals (Whitlow, 2005).

Intoxication by AF in dairy calves is associated with decreased feed intake, body weight loss and death. Lesions in liver include fat accumulation, cellular necrosis, fibrosis, venous occlusion, disorganization of liver lobules, and biliary hyperplasia. Other changes include serum enzyme changes and reductions of vitamin A. Calves up to 6

months of age are most susceptible to the toxic effects of AF but true infections by *Aspergillus* has been linked in heifers and adult cattle to mycotic pneumonia and mycotic abortion (Lynch, 1971).

Adult ruminants have been considered to be less affected by AF because rumen flora could degrade and inactivate mycotoxins (Fink-Gremmels, 2008). However, research does not support this theory completely since biodegradation of AFB₁ in the rumen has shown ranges anywhere from 42% to no apparent degradation in several in vitro experiments (Jouany and Diaz, 2005). Yiannikouris and Jouany (2002) found that concentrations of AF ranging from 0.1 to 10 µg/ml are poorly degraded in the rumen; rumen bacteria are inhibited by AF which may disturb the metabolism and growth of the rumen micro-organisms.

Fink-Gremmels (2008) described that AFB₁ is changed in the rumen to AFL which can be transformed back to AFB₁. According to Coulombe (1993), AFL toxicity is equal to the parent toxin. A significant amount of the AFB₁ that escapes ruminal fermentation is converted in the liver into AFM₁, which is partially excreted into milk (Fink-Gremmels, 2008).

Chronic aflatoxicosis is characterized by poor feed conversion, reduced growth, jaundice, cirrhosis with bile duct proliferation and significant reduction in milk production (Applebaum et al., 1982). Lesions of aflatoxicosis in dairy cattle show liver cell injury, congestion and bleeding, and fatty liver syndrome (pale livers) (Agag, 2004; Fink-Gremmels, 2008).

Edrington et al. (1994) reported that lambs had lower daily feed intake and daily weight gain when 2.5 mg of AF was feed. Semen of buffalo bulls that consumed 2, 3, and

4 grams of rice contaminated with 15.6 mg/kg AF showed a 10 % reduction in live cells and more than 54 % abnormal cell (Hafez et al., 1982).

Aflatoxin in the rumen decreased cellulose digestion and volatile fatty acids production and decreased rumen motility. Cook et al. (1986) fed steers with 0.2, 0.4, 0.6, and 0.8 mg of AF per kilogram of body weight. They observed that the amplitude and frequency of rumen contractions were altered depending on the dose of AF received.

Immunosuppression caused by AF has been connected with disease epidemics in farm animals. A cell-mediated response is particularly sensitive, and phagocytic response is also decreased affecting innate immunity (CAST, 2003). Cordova et al. (2003) looked at the consumption of small quantities of AFB₁ on reproductive performance in dairy cattle and found that immunosuppressant and teratogenic effects of AFB₁ could compromise reproductive parameters. Calving percent in the herd of study was 62%, which is 20 to 30% below of that reported in the literature to be normal and calving intervals was 444 d, 70 d higher than optimal (350 -380 d optimal).

Effects of Aflatoxin on Humans

The impact of food and feed contaminated with mycotoxins, especially AF, is a concern for food organizations worldwide and is expected to receive more attention in coming years (Kamkar, 2005). The problem is more accentuated in developing countries where the food system, economics and infrastructure limits the management of AF contamination (Williams et al., 2004). An investigation by Jonsyn-Ellis (2001) in Sierra Leone showed an average of 96% exposure to AF based on 434 urine samples of school children.

Human exposure to AF is primarily from consumption of contaminated food. However, an inhaled contaminated particle represents another form of AF contact, especially for those workers interacting with grain dust. There are reports where AF was found in the lung of one textile worker, two agriculture workers, a Brazilian engineer who worked with contaminated peanuts and two agriculture employees whom died from pulmonary interstitial fibrosis (Selim et al., 1998).

Aflatoxicosis is the name given to the aflatoxin disease. There are two courses of the disease: acute and chronic. Acute aflatoxicosis results in deaths from hepatic necrosis and liver failure. Chronic aflatoxicosis in humans and animals are related to cancer, immune suppression, hepatocellular carcinoma, reyes syndrome, cirrhosis and kwashiorkor (Stora et al., 1983; Bennett and Klich, 2003).

The symptoms of acute aflatoxicosis disease consists of hemorrhagic necrosis of the liver, bile duct proliferation, edema, lethargy and death (usually in children because human adults are more tolerant to the toxin) (Williams et al., 2004).

Chronic exposure has been related to nutritional and immunotoxicities. Lamplugh and Hendrickse (1982) reported an association between AF and Kwashiorkor. Nelson et al. (1980) and Stora et al. (1983) reported associated AF contamination with Reye's syndrome. Coulombe (1993) suggested that the carcinogenic and mutagenic effects of AF result from the binding of the reactive epoxide with DNA.

There are reports of more than 200,000 deaths annually in the Republic of China due to hepatocellular carcinoma (HCC) related to the consumption of AF (Wang et al., 2002). According to Kamkar (2005) AF is a concern because it may be the contributing agent to human hepatic and extra hepatic carcinogenesis.

Human exposure to AF can be made directly by consuming contaminated feeds like cereals, seeds, fruits, etc., or indirectly by eating food products and subproducts obtained from animals consuming contaminated feeds (Galvano et al., 2005). Milk is an important way to introduce AFM₁ in the human diet. Because milk is the main nutrient for infants and children, the presence of AFM₁ is a concern (Kamkar, 2005). Sadeghi (2009) suggests that exposure to AF prenatally can produce poor neonatal survival and growth retardation.

When dairy products are manufactured from milk contaminated with AFM₁, the toxins are transmitted to the final product because AFM₁ is not degraded when processed into cheese, yogurt, cream and butter. In some kinds of cheeses, AF can be produced by the addition of fungi during the ripening process (Galvano et al., 2005).

Aflatoxin Analysis

Diagnosis of aflatoxicosis in animals is usually performed after other causes have been discarded and, and long after the contaminated feed had been consumed. This reduces the possibilities to obtain samples of feed which caused the problem (CAST, 2003). On the other hand, the analysis of AF is important for enforcing legislature, maintaining the quality of material for feed industry and for estimating occurrence and consumer exposure. Methods of analysis for AF may be classified as fully quantitative, semi-quantitative, or simple yes/no answer, and the requirement for the method will depend on the reason for the analysis (Scudamore, 2005).

According to CAST (2003), major principles were described to ensure a mycotoxin quality assurance program. A good quality assurance program depends on proper implementation of each principle.

Sample Selection and Preparation. Proper sampling is important for decreasing variability in the overall testing. Different commodities have different mycotoxin distribution in the lot. Use of adequate devices like probes to sample bins, rail cars, or piles typically helps to obtain a better sample. When samples are taken from a transferred product, such as conveyor belt or other dynamic lot, small samples can be taken at different intervals of time. At least 100 small samples from the entire lot are recommended (CAST, 2003). Liquids are more homogeneous in the distribution of mycotoxin, but if they contain suspended material, mycotoxins could be unequally distributed (Scudamore, 2005). Sample preparation consists of grinding the sample from a solid phase to facilitate the extraction of mycotoxins (Scudamore, 2005).

Extraction. After grinding, the sample is mixed with a solvent to take out as much mycotoxin as possible. Acetonitrile and methanol are the most common solvents used, and the combination of solvent and water improved recovery better than solvent alone (Scudamore, 2005). Stroka et al. (1999) reported that the ratio between solvent and sample weight affected the final result obtained. Extraction uses two methods: The slow method consists of shaking the sample and solvent from 30 min up to 1-2 hours (CAST, 2003). The other method consists of blending the sample with the solvent for a few minutes. Blending is a more rapid process than shaking, but care must be taken to ensure that the solvent wets all particles (Scudamore, 2005).

Clean-up. A clean-up step to eliminate impurities is recommended but not mandatory for all quantification methods, like ELISA (CAST, 2003). The most common method for clean-up is performed using a solid-phase extraction column which is made of porous silica modified to absorb the impurities or the mycotoxin. One example is the immunoaffinity columns for AFM₁ where solution is passed over the column at a rate of 2-3 ml/min. The precise antibodies are bound to AFM₁, and all other components from the matrix pass through and are washed off using water. The mycotoxin is recovered by washing the column with a solvent (Grosso et al., 2004).

Detection. This final step is classified into two groups according to the ability of the determination assay (see Table 3). Less meticulous methods of detection are called rapid method (CAST, 2003) in which its advantage is to reject or accept materials in rapid commercial decisions (Scudamore, 2005). The other methods of classification constitute quantification of the mycotoxin which is usually more accurate but more expensive (CAST, 2003).

According to Rodriguez et al. (2003) there are advantages of using an ELISA test, such as the reduction in assay time, a simple sample extraction, and specificity for the toxin. On their experiment with milk collected from farms of Leon, Spain, samples run by the ELISA test detected 95.4% positive samples, and the negative samples (4.6%) were run with the HPLC method to detect possible false negatives. All samples were negative to HPLC testing, confirming the efficiency of ELISA essays. However this contrasts with a study reported by Lin et al. (1998) who saw poor results using the ELISA method. Scudamore (2005) published a review of some advantages and disadvantages from methods available for mycotoxin analysis which is summarized in the Table 3.

Table 3. A Review of Methods Available for Mycotoxin Analysis

Method ¹	Advantages	Disadvantages
TLC	<ul style="list-style-type: none"> * Simple, cheap, rapid * Detect most mycotoxins * Sensitive to aflatoxin * Number of samples can run together 	<ul style="list-style-type: none"> * Identification of spots may need confirmation * Insensitive to some toxins * Poor precision * Separation may require 2-dimentional approach
HPTLC	<ul style="list-style-type: none"> * Quantitative when used with densitometry * Number of samples can run together * Improved separation compared with TLC 	<ul style="list-style-type: none"> * Insensitive to some toxins
ELISA	<ul style="list-style-type: none"> * Visual assessment possible * Plate reader assists 	<ul style="list-style-type: none"> * Matrix interference may affect result * Limited to a few solvents * Antibodies may cross react
HPLC	<ul style="list-style-type: none"> * Sensitive, selective * Easy to automate 	<ul style="list-style-type: none"> * Compounds must have UV absorption or fluoresce or require derivatisation
HPLC/MS	<ul style="list-style-type: none"> * Provides high level of confirmation * Multi-detection? * Very sensitive 	<ul style="list-style-type: none"> * Expensive * Specialist expertise required
GC/MS	<ul style="list-style-type: none"> * Provides high level of confirmation * Very sensitive 	<ul style="list-style-type: none"> * Expensive * Specialist expertise required * Compounds must be volatile
CEP	<ul style="list-style-type: none"> * Very low solvent use * Alternative separation technique * Rapid 	<ul style="list-style-type: none"> * Not yet widely evaluated

¹ TLC = Thin layer chromatography; HPTLC = High pressure thin layer chromatography; ELISA = Enzyme linked immunosorbent assay; HPLC = High performance liquid chromatography; HPLC-MS = High performance liquid chromatography with mass spectrometry; GC/MS = Gas column with mass spectrometry; CEP = Capillary electrophoresis.

Fluorometry Analysis

Stokes fluorescence is the re-emission of longer wavelength (lower frequency) photons (energy) by a molecule that has absorbed photons of shorter wavelengths (higher frequency). Both absorption and radiation (emission) of energy are unique characteristics of a particular structure of the molecule during the fluorescence process. Light is absorbed by molecules which cause electrons to become excited to a higher electronic state.

The electrons remain in the excited state for a few seconds then, assuming all of the excess energy was not lost by collisions with other molecules, the electron returns to the ground state. Energy is emitted during the electrons return to their ground state. Emitted light is always a longer wavelength than the absorbed light due to limited energy loss by the molecule prior to emission (Turner Designs, 2008).

Based on the fluorometric technology, VICAM developed AflaTest® a quantitative fluorometric method for the detection of AF in many commodities. This process can be used in a wide variety of locations from the local farm to the government laboratory. It is quick, easy to perform and accurate (Vicam, 1999).

Cucci et al. (2007) felt that in order to ensure reproducibility of fluorometric measurement it is important to have a systematic protocol, because the major disadvantage of the fluorometry method is the variability of quantitative results influenced by very low AFM₁ concentration, sample preparation, presence of impurities, or, air bubbles in the liquid. The same group mentioned that the use of fluorescence

enhancer such as succinyl- β -CD, shows promising result as the most efficient enhancer for AFM₁.

High Pressure Liquid Chromatography

The HPLC uses fluorescence detection, and has been used as the reference technique for detecting AF (Gilbert, 2002; Van Eijkeren et al., 2006). HPLC works well to separate and quantify AF (Wilson et al., 1998). However, this method is complex, laborious, and time-consuming, and it has large requirements for solvents, and needs large investments in costly equipment (Peña et al., 2002). Separation of AFM₁ with HPLC is normally performed using normal or reverse phase methods with fluorimetric detection leading to specificity, high sensitivity, and simplicity of action (Muscarella et al., 2007). A compilation of official methods for AF analysis is presented by Gilbert (2002).

Aflatoxin in Milk

A current serious problem in food hygiene is the occurrence of AF in breast milk, commercial milk and milk products. Considering that milk is a key source of nutrients for humans, this is more important for infants and children whose diet is basically milk. In addition, they are more sensitive to the toxic effects (Galvano et al., 2005).

As the economy improves in developing countries, milk consumption also increases. According to Delgado (2003) a boost of milk consumption in developing countries has been surpassing the growth of milk consumption in the developed countries. The same author estimates that by 2020 milk consumption will be 177 million

metrical tons more than the consumption in 1998. Thus it will elevate the consumption of AF if milk is contaminated.

In North America, milk and other dairy products compose an important part of the diet. According to USDA, the trend in U.S. per capita consumption is downward; in 2001 it was 37 gallons of whole milk per person, eight gallons less than in 1941. But there was an increase in consumption of low fat milk, 15 gallons in 2001 compared with four in 1945. However as the consumption of fluid milk in the U.S. dropped, consumption of cheese increased eight times compared with 1909. This is almost 30 pounds per person and convenience foods such as pizzas are primarily responsible for this increase (Putnam and Allhouse, 2003).

Many studies demonstrate the presence of AF in ingredients used to feed dairy cattle (Yiannikouris and Jouany, 2002; Binder et al., 2007; Driehuis et al., 2008). The conversion of the AF into AFM₁ and its subsequent excretion into milk varies from less than 1 to 3% (Applebaum et al., 1982), to 6% (Galvano et al., 2005) with a median transfer level of 1.7%, disregarding the change in the day to day operation (Jouany and Diaz, 2005).

The presence of AFM₁ in milk is proportional to the contamination of feedstuff. Concentrations of AFM₁ in milk and milk products are controlled in several countries (Boudra et al., 2007). As reported by Van Egmond and Jonker (2004), regulation for AFM₁ in 2002 was established in 98 countries, a growth of 30 %, compared to 1995. Currently the regulations of AFM₁ in milk have one standard applied to the European Union at 0.05 µg/kg and another limit of 0.5 µg/kg applied to the United States and several other European and Latin America countries. The calculation for liver cancer for

both levels is very low, and there is not a noticeable benefit when the limit is reduced from 0.5 to 0.05 µg/kg (Van Egmond and Jonker, 2004). Using the 1.7% carryover rate, Jouany and Diaz (2005) calculated that cows consuming 30 µg/kg AFB₁ in feed easily reach the 0.5 ppb of AFM₁ in milk. The European Union level of 0.05 ppb of AFM₁ will be reached when the diet contains 3 µg/kg of AFB₁.

Various studies have shown that AFM₁ in milk and cheese show a tendency towards lower AFM₁ concentration in the summer months. This is accredited to cows grazing in these months and receiving less concentrated feeds (Van Egmond and Dragacci, 2000; Hussain and Anwar, 2007).

Using the ELISA procedure Ghanem and Orfi (2008) found that local farms in Syria had the highest number of positive samples compared with milk powder which was mostly imported from the European Union, which imposes a strict upper limit for AFM₁. This result is in agreement with CAST (2003) which stated that AFM₁ is more abundant in tropical countries because the temperature and humidity favors the growth of the *Aspergillus* fungi.

The AFM₁ restrictions have used similar analytical methodology (Van Egmond and Dragacci, 2000). The ELISA (Ghanem and Orfi, 2008), TLC (Grosso et al., 2004; Kamkar, 2005), HPLC (Kim et al., 2000; Bognanno et al., 2006) and flourometric (Cucci et al., 2007; Hussain and Anwar, 2007) methods, all reported good, quality results. However, the most important addition to the analysis of AFM₁ was the use of immunoaffinity columns to the clean up step (Van Egmond and Dragacci, 2000).

Aflatoxin Determination in Cheese

As stipulated above, AF exposure in humans is mainly by the ingestion of milk and dairy products (Battacone et al., 2005). If dairy products are manufactured from milk tainted with AFM₁, the toxin is transferred to the final product. It is generally accepted that AFM₁ is unaffected by pasteurization or other milk process such cheese, yogurt, cream, and butter. Cheese may become contaminated with AFM₁ in one of the following two ways; AFM₁ is present in milk during cheese processing due to carryover or AF is produced by flora growing in cheese during maturity. Aflatoxin couples with casein causing about a 2.5 to 3.0 fold higher concentration of AFM₁ in soft cheeses and 3.9 to 5.8 fold higher in hard cheese curd than in milk used to make them (Yousef and Marth, 1989). Battacone et al. (2005) found concentration of AFM₁ in cheese curd twice as high as in milk. Kaniou-Grigoriadou et al. (2005) found low levels of AFM₁ in milk below the tolerance level. However, AFM₁ value increased 4.9 times in cheese curd when feta cheese was made.

Cheese may also be contaminated by the addition of desirable mycoflora during ripening, but its toxicological relevance remains unclear (Galvano et al., 2005). Several studies by Oruc et al. (2006) established that bacterial starter used in cheese production may inactivate the AF during the ripening. Bacteria that produce lactic acid such as bifidobacteria can bind AF and almost eliminate it from the milk.

Storage of cheese may not reduce the concentration of the AFM₁ in the curd. Oruc et al. (2006) determines the stability of AFM₁ during the processing and ripening of a traditional white pickled cheese, a traditional soft Turkish cheese. During a three-month ripening period, they did not observe any reduction in toxin concentrations. Govaris et al.

(2001), working with Telemes cheese, observed after 6 months of storage only a 30% reduction of AFM₁ compared to the initial cheese curd concentration. It should be noted that Talemes cheese is usually consumed after two months of ripening.

Control and Prevention of Aflatoxin M₁

The prevention of AFB₁ contamination in material used to feed dairy cattle has proved to be the most successful system of controlling AFM₁ in human food supply. Wu and Bhatnagar (2008) estimated losses related to AF in the United States were more than \$550 million per year including cost of market rejection and animal health and management. Worldwide the cost is much higher.

There are various attempts to stop the production of aflatoxin under field conditions. Research on developing host resistance of plants shows some potential. Corn breeders have explored the corn genotype for genes related to resistance (Brown et al., 1998). Studies on Midwestern and Mexican corn lines show effective results. The genetic manipulation of pathways in plants to produce antifungal compounds and the use of biocompetitive agents are other areas of experimentation (Brown et al., 1998; Palumbo et al., 2008). Positive crop management practices such as proper fertilization, weed control, irrigation, and insect control reduce AF contamination in corn (Plasencia, 2004).

However, when attempts to stop aflatoxin in the field are not possible or fail, several approaches for post-harvest decontamination can be made, including physical, chemical and biological inactivation (Sinha, 1998). Aflatoxin production post-harvest could be minimized if grains are dried properly before storage, and implementing good management storage practices (CAST, 2003). Mechanical and hand separation has been

used with successful results in nut industries and with corn using flotation and density segregation. The principle is based on removing damaged kernels according to the variation in size, shape, color and visible moldy material (Sinha, 1998).

Chemical inactivation of AF using gaseous ammonia has been successful in animal feeds (Sinha, 1998; Park and Price, 2001). Biological methods, using molds, bacteria, and algae, have reduced AF by competing and excluding toxigenic strains (Sinha, 1998; Yin et al., 2008).

The use of inert absorbents is considered the most promising inactivation treatment. Absorbents like aluminosilicates, clays such as bentonite and zeolite, active carbons, glucomannans and mannanoligosaccharides, aluminosilicates mixed with clays, have been utilized for their ability to sequester AF and reduce their availability for absorption from the gastrointestinal tract (Diaz and Smith, 2005).

Several physical and chemical processes for eliminating or inactivating AFM₁ in milk have been investigated, because degradation of AFM₁ was not observed during industrial processing of milk. Pasteurization, acidification for cream and yogurt production, separation of butter and cheese ripening did not decrease the concentration of AFM₁ to acceptable concentrations (Galvano et al., 2005). Studies of chemicals that could be used for their ability to degrade AFM₁ are limited to those that are permitted as food additives such as sulfites, bisulfites, and hydrogen peroxide (Applebaum and Marth, 1982).

Other processes that have been explored to remove AFM₁ from milk include absorption and radiation. Applebaum and Marth (1982) reported that 5% bentonite in milk adsorbed 89% of AFM₁. In a study of the effects of ultra-violet radiation with and

without hydrogen peroxide, the concentration of AFM₁ was reduced by 3.6–100%, depending on the length of time the milk was exposed to radiation, the volume of treated milk, and the presence of hydrogen peroxide (Yousef and Marth, 1989).

Utilization of materials with chemo-absorption properties has possibilities. Attempts of direct chemi-absorption of AFM₁ in milk were made by Soha et al. (2006) who added bentonite and aluminosilicate at concentrations of 0, 0.05 and 2% to contaminated AFM₁ raw milk. Chemi-absorption compounds significantly reduce milk AFM₁ without affecting milk components.

The chemical and physical treatments described are not readily applicable in the dairy industry, at present, as little is known about the biological safety, or the nutritional value of the treated products. Moreover, the costs of the processes may be considerable and prohibitive for large-scale application (Sinha, 1998). If AFM₁ cannot be easily destroyed or removed from milk, the only way that can be excluded is by eliminating AFB₁ from the animal's diet (Diaz and Smith, 2005)

Developing Methods for Aflatoxin M₁ in Cheese

Advances in developing new methods to study mycotoxins should offer a benefit over existing methods (Scudamore, 2005). New methodology or changes to previous methods need to be supported by comparing results for an existing reference method. After the assay has been evaluated in one or more laboratories, it may go through the process to be established as a new method. Internal validation and collaborative testing provide strong information about method performance and possible problems (CAST, 2003). Methods for validation should take into consideration economical factors as well

as efficiency in time and materials (Gilbert, 2002). Because of frequent contamination of dairy products with AFM₁, the development of fast, accurate and economically viable methods is quite important.

HPLC and TLC methods are the gold standard to determine AFM₁ in milk. There are only a few methods that have been developed for cheese products due to high complexity of their food matrix (Monaci et al., 2007). Cucci et al. (2007) implemented a method to detect of AFM₁ in liquid solutions using a fluorometric process. Once the AFM₁ is extracted from cheese using organic solvent, this method offers several advantages over HPLC. Fluorometric measures of purified extract have been developed because it is a faster method (Stroka and Anklam, 2002) and allows for on-site determination of AF before the final product is made (Sibanda et al., 1999). Fluorometry values are similar to HPLC; however, the results of this method need to be validated with complex matrices such cheese (Stroka and Anklam, 2002).

The ubiquity of mycotoxins causes great economical losses worldwide in the animal production industry. The analysis of AFM₁ is a great concern due to its prevalence in milk and milk products. Consumption of contaminated products can lead to carcinogenic, teratogenic and other detrimental diseases in humans. Of the various methods for mycotoxins detection, HPLC appear to be more sensitive and with automation could be the preferred method. However, equipment is expensive. For this reason it is inaccessible for developing countries or farm analysis. New methods for testing mycotoxins need to be validated that are useful for those sectors.

Interestingly, only minimal or no research exists investigating and validating those new methods for complex matrices such as cheese or dairy products. The present

study attempts to demonstrate that the concentration of AFM₁ in cheese is higher than found in milk and investigate whether fluorometric analysis is comparable to HPLC and could be used to screening and predict AFM₁ from milk products.

MATERIAL AND METHODS

Contaminated Corn with Aflatoxin B₁

Strains of *Aspergillus spp* were planted in flasks containing sterile steam rolled corn. After the analysis, the strain which produced more AFB₁ was grown in a bigger scale. We mixed for 1½ h through a concrete mixer 4 kg of steam-rolled contaminated corn at 400 µg/kg to with 36 kg of steam rolled AF free corn to obtain corn with 40 µg/kg. Bags containing 1.6 kg and 0.125 kg, respectively, were weighed, marked, and stored until their use for feeding cows in high and low treatment groups.

Animals

Four Holstein cows in late lactation (>220 DIM), were used from the USU Caine Dairy Teaching and Research Center herd. All animal were allowed one week acclimation period to the tie stall facility. The cows were milked twice daily at 0530 and 1800 h and fed a TMR diet consisted of (corn silage (35%), alfalfa hay (25%), and energy-protein supplement (40%)) for ad libitum intake. Ration was formulated to meet the nutrient requirement of cows producing 30 ± 5 kg of 3.5% FCM/d according to NRC (2001) recommendation. Composition of the diet for each treatment is given in Table 4. Cows were feed twice daily (0600 h and 1730 h) ad libitum. The cows were divided in two groups. One group (high group) of two cows received 1.6 kg of contaminated corn and the other group (low group) received 0.125 kg of contaminated corn mixed with 1.475 kg of AF free steam rolled corn (to equal the matrix intake of the high treatment group) immediately after morning milking. To ensure cows consumed the contaminated grain, corn was put in a plastic container before the TMR feeding and consumption was

monitored. Fresh water was provided ad libitum. Animal care and procedures were approved and conducted under established standards of the Utah State University Institutional Animal Care and Use Committee.

Table 4. Ingredients and Chemical Composition of Diet

Composition ¹	
Ingredient	g/100 g of DM
Alfalfa hay	25.69
Corn silage	35.42
Steam rolled corn	12.63
Corn, hominy	9.55
Canola meal mechanically extracted	3.69
Whole-linted cottonseed	5.92
Soybean meal expeller	2.04
Blood meal, ring dried	1.00
Molasses, sugar beet	2.50
Minerals & vitamin mix	1.56
Chemical	
NE _L , Mcal/kg of DM	1.44
CP	16.6
NDF	32.8
ADF	21.9

¹Extra 1.6 kg of steam rolled corn were fed to each cow. High AF treatment were given 1.6 kg of contaminated corn and low treatment were given 0.125 kg of contaminated steam rolled corn added to 1.475 kg of free AF steam rolled corn

Milk Collection and Processing

Cows were milked twice daily (0530 h and 1700 h) for 10 d each into vacuum-sealed milking cans. Milk used for processing into cheese was taken from the 5 and 7 d morning milking after AFB₁ addition in the diet. Milk from the high and low AF treatments was collected in 15 L buckets on the dairy farm. Milk was pooled and brought to the Richardson Dairy Products Laboratory at Utah State University (USU) and

transferred into a 15 L steel bucket. Milk not used for cheese production was disposed into the drain on the farm.

Cheese Production

The cheese production was during April of 2007. Two sets of cheeses, Cheddar and Fresco, with two AFM₁ concentrations, high and low were made for each concentration of AFM₁ in milk. The following are some characteristic describing each cheese batch. Cheddar cheese is hard-cooked, and long-matured cheese, it has a firm texture, pale yellow to orange color with different intensity of their sharp flavor going from mild to extra-sharp. Composition of Cheddar cheese consists of water content of 30 to 32%, 34 % protein and 29% fats. On other hand, Fresco cheese is bland high moisture, usually lower in fat and sodium than other cheeses. Composition of Fresco cheese is 59% water, 19% protein and 15% lipid content (Dr. Nicoletta Fuca, personal communication).

The 15 L buckets were placed in a small steam jacketed vat. Adequate water was added to the vat, without tipping over the containers. The milk buckets were heated to 65°C for Cheddar cheese and 73°C for Fresco cheese by heating the water and stirring the milk. When temperatures were reached, the water was drained and replaced with cold water. The milk buckets were cooled to the set point for cheese of 31°C for Cheddar and 33°C for Fresco cheese. This provided a pseudo-pasteurization heat treatment to facilitate cheese making. Twenty liters of milk from each treatment (AF high and AF low) were used to make each type of cheese.

Making of Cheddar Cheese

Milk from both treatments of AF (high vs. low) was converted into Cheddar cheese using a stirred curd method based upon typical manufacturing procedures at Richardson Dairy Products Laboratory at USU. Temperature for cheese production was set at 31°C. After reaching that temperature 3 g of DVS 850 standard culture, 1.5 ml of CaCl₂, 1 ml of annatto colorant and 1 ml of coagulant were added. This was stirred for two min. and then left undisturbed to ripen until ready to cut. Curds were cut and left undisturbed for 5 min. The curds were stirred and slowly cooked until the temperature reached 49°C. Whey was then drained, and curds were stirred vigorously to dry the curd. Salt was added (33 g) when the pH reached 5.3. Curds were pressed overnight in tubes at 20 psi to yield approximately 0.5 kg blocks of Cheddar cheese. Blocks were cut into 0.2 kg samples labeled, vacuum packaged and stored at 7°C for 15 d.

Making of Fresco Cheese

Milk from cows administered high or low AF was converted into Fresco cheese based upon typical manufacturing procedures at Richardson Dairy Products Laboratory at USU as followed; milk temperature was set at 34°C, and then 4 g of TC-20C *Strep. thermophilus* culture was added and the milk left undisturbed for 15 min, then, 1 ml of DS chymosin (aspartic acid enzyme) was added as rennet. Curds were cut after 30 min into 1.5 cm cubes and 8 L of the whey were drained and replaced with hot water. The curds were stirred for 30 min until they reached pH 5.7. Salt was added (70 g). Curds were filled into cloth lines tubes and pressed for 2 h at 5 psi. Blocks were cut into 0.2 kg samples, labeled, vacuum packaged, and stored at 3°C for 15 d.

Detection and Quantification of Aflatoxin M₁ in Cheese

Chemicals, Standards and Laboratory. The fluorometer calibration standards Aflatest-MTM, the immunoaffinity columns AflaM1TM and Developer were purchased from VICAM (Watertown, MA). All organic solvents were obtained from Fisher Scientific (Pittsburgh, PA). Deionization water used in the experiments was processed by a Barnstead water purification system (Barnstead PCS, Dubuque, IA). Microfiber filters (0.45μm) were obtained from Fisher Scientific. All other materials such as disposable pipets, filter paper, microfilters, disposable cubets and syringes were purchased at the chemist store at USU. Extraction and analysis of AF was made in a room protected from daylight and vials were of amber color or protected with foil paper, because AF are degraded by UV light.

Extraction of Aflatoxin M₁ from Cheese. Aflatoxin M₁ was extracted from cheese using the method described by Dragacci et al. (1995) with some modification. Cheese was brought to room temperature and then cut into small pieces and minced thoroughly. On a clean table minced cheese was distributed to form a square. Diagonals were traced on the square, and two of the opposite triangle sides were discarded. The remaining triangles went through the same process until the sample was around 20 g. A 10 g subsample of cheese was blended with 10 g of celite (Diatomaceous earth powder) 545 (Fisher Scientific Inc.), and 80 ml dichloromethane, for two min. at high speed (Waring blender, Model No 35BL 64, Merck, Poole, UK). After washing further with 40 ml dichloromethane, the mixture was filtered (filter paper #1 circles 24.0 cm; Whatman) and pressed to release maximum amount of filtrate. Each cheese was analyzed in triplicate.

The filtrate was evaporated at 40°C on a rotary flash evaporator (Buchler Instruments, Fort Lee, NJ) with a precision stainless steel water bath Model 183. The residue was dissolved in 1 ml methanol, 30 ml water (1:30, v/v) and 50 ml n-hexane, and then transferred to a separating funnel. The lower (water phase) layer was collected and then the hexane phase was washed twice with 10 ml water. The water phase was collected after each wash.

Bottles of the collected water fraction were wrapped in paper foil to avoid light contact and stored at 4° C until processed. The aqueous phase was passed through an AFM₁-HPLC immunoaffinity column (AFM₁ monoclonal antibody-based affinity chromatography of VICAM, Watertown, MA) at a rate of 2-3 ml/min. This process was performed in a solid phase extraction manifold (SPE Manifold, Waters, Watertown, MA). As the sample passed through the column, the antibodies selectively bound with AFM₁ forming a complex. Then the immunoaffinity column was washed twice with 10 ml of distilled water to eliminate impurities. The toxin was eluted using 2 ml methanol. Once the AF samples were eluted, the 2 ml liquid was divided into 0.5 ml vials for HPLC measurement and 1 ml glass tubes for fluorometric measurement.

Aflatoxin M₁ Determination

Fluorometric Determination. The analyses were performed by a VICAM Series-4ex Fluorometer optical system (VICAM, Watertown, MA) equipped with a high intensity pulsed Xenon lamp, together with selected fluorescence excitation and emission filters. Linear calibration of the instrument was developed by subsequently reading three quinine sulfate dehydrate reference standards (Aflatest-^{MTM}; VICAM). The first and the

second vials were used to set the higher and lower calibration points and the last vial was used to verify the instrument linearity, as referred to in the VICAM manual.

The reading range was fixed at 0.01 – 22 $\mu\text{g}/\text{kg}$ for AFM₁, with a precision of 0.5% and accuracy of 1%, as reported in the VICAM manual. Two ml of reagent (1ml methanol + 1mL developer) and 2mL of distilled water were consecutively read as blank samples. In the fluorometric method the AFM₁ elution was mixed with developer solution. Aflatoxin enhancer (VICAM developer), should be prepared daily. It is not recommended to use if color is lost or after six hours have elapsed. According to manufacture one part of concentrate developer is mixed with nine parts of distilled/deionized water. Then 1 ml of enhancer solution is used with 1 ml of eluted AFM₁ and read in the fluorometer. Analyses were performed at the same time as the HPLC was injected with the same sample (approximately every ten minutes). All cheese samples were run in triplicate.

High Pressure Liquid Chromatography. The HPLC analysis was performed using a Waters 2695 Separations Module equipped with a Waters 2996 PAD and a Waters 2475 multi λ fluorescence detector (Waters, Watertown, MA). The system and data acquisition were controlled by, EmpowerTM Chromatography PC software for the PDA detector (Waters, Watertown, MA). The AFM₁ was separated with a Symmetry[®] C18 LiChospher 100 column with 3.5 μm particle size (150 x 4.6 mm I.D.) at room temperature, with acetonitrile/water (25:75 v/v) as the mobile phase (flow rate one ml/min). The fluorescence detector was set at 365 nm excitation and 440 nm emission wavelengths. Standard AFM₁ (Sigma Chemicals Co., St. Louis, MO) was used to prepare a series of working standards of 12, 120, 240, 360 and 480 $\mu\text{g}/\text{AFM}_1/\text{L}$. Calibration

curves were arranged by plotting the peak area for each standard against the quantity of AFM₁ injected. All analyses were done in triplicate, injecting 100 µl into the HPLC.

Stability of the Method

Stability of the method was assessed by measurement AFM₁ from a cheese sample on four different days. The same extraction and analyses as above was performed.

Statistical Analyses

The design of the experiment was a 2-way Fixed Factor Analysis. Factor A concentration of AF (high and low) and factor B cheese type (fresco and cheddar) with 30 (cheese) repetitions. Data were analyzed using the PROC MIXED function in SAS (SAS, 2009). The dependent variables were methods of analysis and the independent variable was cheese type (Cheddar and Fresco), treatment (high and low AFM₁) and an interaction term. Least squares means were computed and means differences determinate by the Tukey's multiple means comparison with significance at P < 0.05. Pearson's correlation was run between the two analysis methods and the PROC REG function in SAS used to determine the linear relationship between the two analysis methods.

A linear relationship was computed and the correlation is represented by the equation (y = mx + b), where m = coefficient, x = fluorometer or hplc value and, b = intercept.

RESULTS AND DISCUSSION

Carryover of Aflatoxin M₁ Feeding Trail

A sample of milk for each treatment level was taken two days prior to cheese preparation and analyzed for AFM₁ in our laboratory by the HPLC technique (Dragacci et al., 1995). Another sample was sent to an external laboratory. Table 5 summarizes the concentration obtained for those samples.

Table 5. Concentration of Aflatoxin M₁ in Milk Prior to Cheese Preparation. Evaluation was made April 16, 2007 by the USU lab and February 12, 2009 by the external lab.

Treatment ¹	USU lab, µg/kg	External lab, µg/kg
High	0.8	0.7
Low	0.03	Trace

¹Milk was pooled for both cows into one treatment

The experiment was not set up to measure AFM₁ carryover. However, based on the amount of grain fed to each group and the concentration of AFB₁ in grain, we retrospectively calculated the possible carryover. We calculated that the carryover of AFM₁ from the high treatments was approximately 1.3% and carryover for the low treatment of AFM₁ was 0.75%. The high AF group receive 1.6 kg of corn daily containing 40 µg/kg of AFB₁ ($1.6 \text{ kg} \times 40 \text{ µg/kg} = 64 \text{ µg/kg}$ $\times 1.3\% \text{ carryover} = 0.8 \text{ µg/kg}$ of AFM₁). The low group received the same amount of corn but only 0.125 kg were from contaminated corn at 40 µg/kg of AFB₁ ($0.125 \text{ kg} \times 40 \text{ µg/kg} = 5 \text{ µg/kg}$ $\times 0.75\% = 0.03 \text{ µg/kg}$). These concentrations represent the two universal restrictions for AFM₁, 0.03 µg/kg is below European regulatory value (0.05 µg/kg) and 0.8 µg/kg is over

American restriction (0.5 µg/kg). Those values are in accordance with Applebaum et al. (1982) and Jouany and Diaz (2005).

Detection of Aflatoxin M₁ in Cheese

To our knowledge, this is the first report using a fluorometric procedure to analyze AFM₁ in cheese. The two types of cheeses were chosen to represent a moist, soft cheese and a dry, hard cheese in order to evaluate whether the cheese type had any impact on AFM₁ analysis.

The main objective was to compare HPLC vs. fluorometry measurement of AFM₁ in cheese. The presence of AFM₁ was detected in all samples (100%) by both methods. Table 6 shows the statistical analysis for the cheese type, treatment, and analysis method.

Table 6. Means of Aflatoxin M₁ for High Performance Liquid Chromatography and Fluorometry Method From Different Cheese Types and Aflatoxin M₁ Fed Concentrations

Cheese	Treatment	Method	Mean	SD	Range	CV(%)
Cheddar	High	HPLC	1.03 ^a	0.41	0.22 - 1.98	39.52
		Fluorometry	1.36 ^a	1.48	0.29 - 8.06	108.9
	Low	HPLC	0.03 ^b	0.01	0.01 - 0.05	34.09
		Fluorometry	0.02 ^b	0.02	0.01 - 0.10	86.93
Fresco	High	HPLC	0.93 ^a	0.36	0.32 - 2.06	38.86
		Fluorometry	0.95 ^a	0.27	0.40 - 1.56	28.04
	Low	HPLC	0.04 ^b	0.03	0.01 - 0.19	81.94
		Fluorometry	0.05 ^b	0.02	0.01 - 0.08	44.60

^{a-b} Means within a column with different superscripts differ (P<0.01)

There were no significant differences between cheese type (P<0.05). Amounts of AFM₁ in feed resulted in difference of AFM₁ in cheese regardless of its type (P<0.01). Interactions between concentration and cheese sample or cheese type were not statistical

significant at ($P>0.05$). There were no differences between methods of detection with treatment ($P>0.05$).

Yousef and Marth (1989) reported that the concentration of AFM₁ in cheese was about 3.0 (2.5 – 3.5) and 4.8 (3.9 – 5.8) fold higher for soft and hard cheese than that found in milk, respectively. In this study AFM₁ concentration in both cheeses was higher than they found in milk, except for low Cheddar cheese batch, but not as high as reported above by Yousef and Marth.

Carryover values of the AFM₁ (Table 7) were over 100% for the high concentration of AFM₁ in Cheddar and Fresco and for low concentration in Fresco cheese, respectively (average of 132%). Cheddar cheese low concentration of AFM₁ had lower carryover rate (89%) than other cheese batches. This may be due to limited casein proteolysis. Gürses et al. (2004) suggested that proteolysis of casein may increase the recovery of AFM₁ from cheese.

Table 7. Mean Carryover of Aflatoxin M₁ in Cheese as Compared to Milk Content

	Cheese ¹			
	Cheddar		Fresco	
	High	Low	High	Low
HPLC	128	97	115	133
Fluorometric	163	80	119	133

¹Values are expressed in %

Based on the higher carryover for AFM₁ in cheese (Table 7), we also conclude that AF is heat stable. These results are in concordance with Barbiroli et al. (2007) who reported that AF was stable at temperatures used during cheese production.

Another difference found between methods of cheese production was the amount and kind of starter bacteria used to initiate the ripening process. In fresh cheese, 4 g of *Thermophilus* bacteria (TS-20CS) was used. In Cheddar cheese, 3 g of a starter culture (DVS 850) was used. Sinha (1998) reported that bacteria and fungus potentially degraded or modified AF.

A linear regression model was derived from data in Table 6. HPLC ($\mu\text{g/kg}$) = $0.303 + 0.344$ (fluorometer value), $r^2 = 0.535$ for both methods, $t = 8.04$ ($P < 0.01$). The regression analyses between HPLC and fluorometry (Figure 1 and Figure 2) shows that fluorometry is a good predictor to screen AFM₁ concentration in cheese.

Stability of the Analytical Method

Stability is a validated quantitative analytical procedure that can detect the changes with time of the active ingredient (CIPAC, 1999). Results of the stability for fluorometry and HPLC are summarized in Table 8. The fluorometry method showed good CV(%) for day one then increased by day 2 to slightly over 20%, and then increases again on day 3. For HPLC, the CV(%) stayed very close together for all 4 d. A possible recommendation for fluorometry is to use fresh samples during each determination to avoid variability on the detection of AFM₁ concentration in cheese.

Table 8. Stability of the Analytical Method for Aflatoxin M₁ Detected by Fluorometry and HPLC

Day	Fluor				HPLC			
	1	2	3	4	1	2	3	4
Mean	0.09	0.11	0.09	0.09	1.24	1.28	1.08	1.28
SD	0.01	0.02	0.02	0.03	0.03	0.02	0.02	0.07
CV%	8.27	20.11	29.21	28.24	2.10	1.68	1.72	2.81




Figure 1. Relationship of Aflatoxin M₁ High and Low Concentration Between HPLC and Fluorometry Method From Cheddar Cheese




Figure 2. Relationship of Aflatoxin M₁ High and Low Concentration Between HPLC and Fluorometry From Fresco Cheese

Fluorometry Cross-Validation

Several properties are required to make a laboratory assay suitable for assimilation into commercial protocol, properties such as sensitivity, specificity, and accuracy (Shah, 2007). A cross validation is the comparison of two or more methods to generate data for the same study. In our study we use HPLC as a reference method and fluorometry as the method to be validated. We follow the criteria published by CIPAC 3870 (CIPAC, 1999).

Estimation of the Precision. Data obtained from calibration and readings by AFM₁ blank dilution were correct and indicated that repeatability was acceptable.

Demonstration of Accuracy. In the present study, AFM₁ used to calibrate the fluorometer was purchased with certification of authenticity and handled as recommended. To avoid the interference from excipients, AFM₁ was extracted the same way for both methods. The results were found to be not significantly different ($P<0.01$) between fluorometry and HPLC. In all cases we found a lack of interference.

Comparison of the Results Obtained by the Two Methods. Each cheese batch was compared; results were the same for both methods of analysis. The fluorometry method is validated for the detection of AFM₁ in cheese.

According to Gilbert (2002), methods for validation should take into consideration economical factors as well as efficiency in time and materials. We evaluate the cost of the fluorometry compared to the HPLC method, respectively. Table 9 shows comparison between those two methods.

Comparing the cost of the equipment between the fluorometer and HPLC, the HPLC is almost 20-fold greater than the fluorometer. Considering the variable time, the

HPLC took longer per sample. Although an automatic injection was used, time average for each sample is 10 min. Taking into consideration the variable cost of analysis, material for running the HPLC include HPLC quality solvents which are more expensive than regular solvents.

Table 9. Comparison of Variables From Fluorometry Against High Pressure Liquid Chromatography

	Methods	
	HPLC	Fluorometry
Equipment cost	\$78,000.00	\$4,000.00
Time ¹	20 ²	2
Difficult	Very difficult ³	Easy ⁴
Variable cost ⁵	\$480.00	\$273.00

¹Time calculated by 120 samples

²Did not include time to warm the machine, washed columns time, respectively

³Include standards calibration and curves determination

⁴Print read-out containing the results in ppb ($\mu\text{g}/\text{kg}$)

⁵Did not include labor fees

CONCLUSIONS

The reduced time to run a sample, the lower cost of the fluorometer machine and the lower variable costs of analysis make the fluorometry method more useful than the HPLC method for screening samples. A printable readout with the amount of AFM₁ can be obtained from fluorometer and eliminate some of variability due to determination at the beginning and end of HPLC curves.

The fluorometric method is simple and reliable. Although there were slight numerical differences between the HPLC and fluorometry methods in detecting AFM₁ in different types of cheeses, those were not statistically different.

This study also indicates that the AFM₁ immunoaffinity column developed for milk, plus the different solvents can be used for AFM₁ detection in cheese. Carryover rate, compared to original milk content, of the AFM₁ in cheeses differs from those previously reported in the literature. Those differences may be attributed to cheese preparation. It will be interesting to compare the protocol used in this study at other cheese factories.

REFERENCES

Agag, B. I. 2004. Mycotoxins in foods and feeds 1-Aflatoxins. *Ass. Univ. Bull. Environ. Res.* 7(1):176-206.

Applebaum, R. S., R. E. Brackett, D. W. Wiseman, and E. H. Marth. 1982. Responses of dairy cows to dietary aflatoxin: Feed intake and yield, toxin content, and quality of milk of cows treated with pure and impure aflatoxin. *J. Dairy Sci.* 65(8):1503-1508.

Applebaum, R. S. and E. H. Marth. 1982. Use of sulphite or bentonite to eliminate aflatoxin M1 from naturally contaminated raw whole milk. *Zeitschrift für Lebensmitteluntersuchung und-Forschung A.* 174(4):303-305.

Barbiroli, A., F. Bonomi, S. Benedetti, S. Mannino, L. Monti, T. Cattaneo, and S. Iametti. 2007. Binding of aflatoxin M₁ to different protein fractions in ovine and caprine milk. *J. Dairy Sci.* 90(2):532-540.

Battacone, G., A. Nudda, M. Palomba, M. Pascale, P. Nicolussi, and G. Pulina. 2005. Transfer of aflatoxin B1 from feed to milk and from milk to curd and whey in dairy sheep fed artificially contaminated concentrates. *J. Dairy Sci.* 88(9):3063-3069.

Bennett, J. W., and M. Klich. 2003. Mycotoxins. *Clin. Microbiol. Rev.* 16(3):497-516.

Binder, E. M., L. M. Tan, L. J. Chin, J. Handl, and J. Richard. 2007. Worldwide occurrence of mycotoxins in commodities feeds and feed ingredients. *Anim. Feed Sci. Technol.* 137(3-4):265-282.

Bognanno, M., L. La Fauci, A. Ritieni, A. Tafuri, A. De Lorenzo, P. Micari, L. Di Renzo, S. Ciappellano, V. Sarullo, and F. Galvano. 2006. Survey of the occurrence of aflatoxin M1 in ovine milk by HPLC and its confirmation by MS. *Molecular Nutrition & Food Res.* 50(3):300-305.

Boudra, H., J. Barnouin, S. Dragacci, and D. P. Morgavi. 2007. Aflatoxin M1 and ochratoxin A in raw bulk milk from French dairy herds. *J. Dairy Sci.* 90(7):3197-3201.

Brown, R. L., D. Bhatnagar, T. E. Cleveland, and J. W. Cary. 1998. Recent Advances in Preharvest Prevention of Mycotoxin Contamination. Pages 351-380 in *Mycotoxins in Agriculture and Food Safety*. K. K. Sinha and D. Bhatnagar, eds. Marcel Dekker, Inc., New York, NY.

Carnaghan, R. B. A., and K. Sargeant. 1961. The toxic of certain ground nut meals to poultry. *Vet. Rec.* 73 (29): 726-727.

CAST. 2003. Mycotoxins: Risk in plants, animals and human systems. Council for Agriculture Science and Technology, Ames, IA.

CIPAC. 1999. Guidelines on method validation to be performed in support of analytical methods for agrochemical formulation. Collaborative international pesticides analytical council. No 4105. Black Bear Press, Cambridge.

Coffey R., E. Cummins, and S. Ward. 2009. Exposure assessment of mycotoxins in dairy milk. *Food Control.* 20(3):239-249.

Cook, W. O., J. L. Richard, G. D. Osweiler, and D. W. Trampel. 1986. Clinical and pathological changes in acute bovine aflatoxicoses: Rumen motility and tissue and fluid concentrations of aflatoxin B1 and M1. *Am. J. Vet. Res.* 47:1817.

Cordova, I. A., O. V. Dorantes, H. R. Escobar, M. G. Iñiguez, M. C. R. Muñoz , B. S. D. Peña, and M. G. A. Trampe. 2003. Efecto de aflatoxinas sobre parametros reproductivos en vacas Holstein. *Med. Vet.* 20(4): 41-44.

Coulombe, R.A. 1993. Biological actions of mycotoxins. *J. Dairy Sci.* 76. 880-891.

Cucci, C., A. G. Mignani, C. Dall'Asta, R. Pela, and A. Dossena. 2007. A portable fluorometer for the rapid screening of M₁ aflatoxin. *Sensors and Actuators B: Chemical* 126(2):467-472.

Delgado, C. L. 2003. Rising consumption of meat and milk in developing countries has created a new food revolution. *J. Nutr.* 133(11):3907-3910

Diaz, D., W. Hagler, J. Blackwelder, J. Eve, B. Hopkins, K. Anderson, F. Jones, and L. Whitlow. 2004. Aflatoxin Binders II: Reduction of aflatoxin M1 in milk by sequestering agents of cows consuming aflatoxin in feed. *Mycopathologia* 157(2):233-241.

Diaz, D., and T. K. Smith. 2005. Mycotoxin sequestering agents: Practical tools for the neutralization of mycotoxins. Pages 323-339 in *The Mycotoxin Blue Book*. D. Diaz, ed. Nottingham University Press, Nottingham, UK.

Draggacci, S. E. Gleizes, J. M Fremy, and A. A. G Candlish. 1995. Use of immunoaffinity chromatography as a purification step for the determination of aflatomin M1 in cheeses. *Food Additives and Contaminants* 12 (1): 59-65.

Driehuis, F., M. C. Spanjer, J. M. Scholten, and M. C. te Giffel. 2008. Occurrence of mycotoxins in feedstuffs of dairy cows and estimation of total dietary intakes. *J. Dairy Sci.* 91(11):4261-4271.

Eaton, D. L., and E. P. Gallagher. 1994. Mechanisms of aflatoxin carcinogenesis. *Annual Review of Pharmacol. and Toxicol.* 34(1):135-172.

Edrington, T. S., R. B. Harvey, and L. F. Kubena. 1994. Effect of aflatoxin in growing lambs fed ruminally degradable or escape protein sources. *J. Anim. Sci.* 72(5):1274-1281.

Elgerbi, A. M., K. E. Aidoo, A. A. G. Candlish, and R. F. Tester. 2004. Occurrence of aflatoxin M1 in randomly selected North African milk and cheese samples. *Food Additives & Contaminants* 21(6):592-597.

FAO. 2004. Worldwide regulations for mycotoxins in food and feed in 2003. FAO, Rome.

Fink-Gremmels, J. 2008. The role of mycotoxins in the health and performance of dairy cows. *The Veterinary J.* 176(1):84-92.

Flannigan, B. 1991. Mycotoxins. Pages 226-257 in *Toxic Substances in Crop Plants*. J. P. F. D'Mello, C. M. Duffus, and J. H. Duffus, eds. The Royal Society of Chemistry, Science Park, Cambridge.

Galvano, F., A. Ritieni, G. Piva, and A. Pietri. 2005. Mycotoxins in the human food chain. Pages 187-224 in *The Mycotoxin Blue Book*. D. Diaz, ed. Nottingham University Press, Nottingham, UK.

Gervais, P., and M. Bensoussan. 1994. Solid-state fermentation of the genus *Aspergillus*. Pages 101-140 in *Aspergillus*. J. E. Smith, ed. Plenum Press, New York, NY.

Ghanem, I., and M. Orfi. 2009. Aflatoxin M1 in raw, pasteurized and powdered milk available in the Syrian market. *J. Food Cont.* 20(6): 603-605.

Gilbert, J. 2002. Validation of analytical methods for determining mycotoxins in foodstuffs. *Trends in Anal. Chem.* 21(6-7):468-485.

Gimeno, A., and L. M. Martins. 2006. Mycotoxins and Mycotoxicosis in Animals and Humans. Special Nutrients, Inc., Mexico City, Mexico.

Govaris, A., V. Roussi, P. A. Koidis, and N. A. Botsoglou. 2001. Distribution and stability of aflatoxin M1 during processing, ripening and storage of telemes cheese. *Food Additives & Contaminants* 18(5):437-443.

Grosso, F., J. M. Fremy, S. Bevis, and S. Dragacci. 2004. Joint IDF-IUPAC-IAEA(FAO) interlaboratory validation for determining aflatoxin M1 in milk by using immunoaffinity clean-up before thin-layer chromatography. *Food Additives & Contaminants* 21(4):348-357.

Gürses, M., A. Erdoğan, and B. Çetin. 2004. Occurrence of aflatoxin M1 in some cheese types sold in Erzurum, Turkey. *Turk. J. Vet. Anim. Sci.* 28(3):527-530.

Hafez, A. H., S. E. Megalla, and A. A. Mahmed. 1982. Aflatoxin and aflatoxicosis. *Mycopathologia* 77(3):141-144.

Hussain, I., and J. Anwar. 2007. A study on contamination of aflatoxin M1 in raw milk in the Punjab province of Pakistan. *Food Control* 19(4):393-395.

IACR. 1997. Some Naturally Occurring Substances: Food Items and Constituents, Heterocyclic Aromatic Amines and Mycotoxins. Vol. 56. World Health Organization. Lyon, France.

Jonsyn-Ellis, F. 2001. Seasonal variation in exposure frequency and concentration levels of aflatoxins and ochratoxins in urine samples of boys and girls. *Mycopathologia* 152(1):35-40.

Jouany, J.-P., and D. Diaz. 2005. Effects of mycotoxins in ruminants. Pages 295-321 in The Mycotoxin Blue Book. D. Diaz, ed. Nottingham University Press, Nottingham, UK.

Kabak, B., A.D. Dobson, and I. Var. 2006. Strategies to prevent mycotoxin contamination of food and animal feed: A review. *Crit. Rev. Food Sci. Nutr.* 46(8): 593-619.

Kamkar, A. 2005. A study on the occurrence of aflatoxin M1 in raw milk produced in Sarab city of Iran. *Food Control* 16(7):593-599.

Kaniou-Grigoriadou, I., A. Eleftheriadou, T. Mouratidou, and P. Katikou. 2005. Determination of aflatoxin M1 in ewe's milk samples and the produced curd and Feta cheese. *Food Control* 16(3):257-261.

Kim, E. K., D. H. Shon, D. Ryu, J. W. Park, H. J. Hwang, and Y. B. Kim. 2000. Occurrence of aflatoxin M1 in Korean dairy products determined by ELISA and HPLC. *J. Food Additives & Contaminants* 17(1):59-64.

Kozakiewicz, Z., and D. Smith. 1994. Physiology of aspergillus. Pages 23-38 in Aspergillus. J. E. Smith, ed. Plenum Press, New York, NY.

Krska, R. 1999. Mycotoxins of growing interest: Zearalenone1. Pages 1-12 in Third Joint FAO/WHO/UNEP International Conference on Mycotoxins, Tunis, Tunisia.

Lamplugh, S. M., and R. G. Hendrickse. 1982. Aflatoxins in the livers of children with Kwashiorkor. *Ann. Tropical Pediatrics* 2:101-104.

Lin, L., J. Zhang, P. Wang, Y. Wang, and J. Chen. 1998. Thin-layer chromatography of mycotoxins and comparison with other chromatographic methods. *J. Chromatogr. A.* 815(1):3-20.

Lynch, G.P. 1971. Mycotoxins in feed and their effect on dairy cattle. *J. Dairy Sci.* 55(9):1243-1255.

Monaci, L., R. Vatinno, and G. De Benedetto. 2007. Fast detection of cyclopiazonic acid in cheese using Fourier transform mid-infrared ATR spectroscopy. *European Food Res. and Techno.* 225(3):585-588.

Muscarella, M., S. L. Magro, C. Palermo, and D. Centonze. 2007. Validation according to European Commission Decision 2002/657/EC of a confirmatory method for aflatoxin M1 in milk based on immunoaffinity columns and HPLC with fluorescence detection. *Anal. Chem.* 594(2):257-264.

National Research Council. 2001. Nutrient Requirements of Dairy Cattle. 7th ed. Natl. Acad. Sci., Washington, DC.

Nelson, D. B., R. Kimbrough, P. S. Landrigan, A. W. Hayes, G. C. Yang, J. Benanides, D. M. Morens, D. Morse, M. Pollack, K. E. Powell, and J. Z. Sullivan-Bolyai. 1980. Aflatoxin and Reye's syndrome: A case control study. *Pediatrics.* 66(6):865-869.

Nuryono, N., A. Agus, S. Wedhastri, Y. B. Maryudani, F. M. C. Sigit Setyabudi, J. Böhm, and E. Razzazi-Fazeli. 2009. A limited survey of aflatoxin M1 in milk from Indonesia by ELISA. *Food Control. J. Food Cont.* 20(8): 721-724.

Oruc, H. H., R. Cibik, E. Yilmaz, and O. Kalkanli. 2006. Distribution and stability of aflatoxin M1 during processing and ripening of traditional white pickled cheese. *Food Additives & Contaminants* 23(2):190-195.

Palumbo, J. D., T. L. O'Keeffe, and H. K. Abbas. 2008. Microbial interactions with mycotoxicogenic fungi and mycotoxins. *Toxin Reviews* 27(3):261 - 285.

Park, D. L. 1993. Controlling aflatoxin in food and feed. *Food Technol.* 47(10):92.

Park, D. L., and W. D. Price. 2001. Reduction of aflatoxin hazards using ammoniation. *Rev. Environ. Contamin. Toxicol.* 171:139-175.

Peña, R., M. C. Alcaraz, L. Arce, A. Ríos, and M. Valcárcel. 2002. Screening of aflatoxins in feed samples using a flow system coupled to capillary electrophoresis. *J. Chromatogr. A.* 967(2):303-314.

Plasencia, J. 2004. Aflatoxins in Maize: A Mexican perspective. *Toxin Reviews* 23(2):155 - 177.

Prado, G., M. S. Oliveira, M. L. Pereira, F. M. Abrantes, L. G. Santos, and T. Veloso. 2000. Aflatoxin M1 in samples of "minas" cheese commercialized in the city of Belo Horizonte - Minas Gerais/Brazil. *Ciência e Tecnologia de Alimentos.* 20:398-400.

Putnam, J., and J. Allshouse. 2003. Trends in U.S. per capita consumption of dairy products, 1909 to 2001. *Amber Waves* 1(3): 12-13.

Richard, J. L. 2007. Some major mycotoxins and their mycotoxicoses: An overview. *Int. J. Food Microbiol.* 119(1-2):3-10.

Riley, R. T., and J. Pestka. 2005. Mycotoxin: metabolism, mechanism and biochemical markers. Pages 279-294 in *The Mycotoxin Blue Book*. D. Diaz, ed. Nottingham University Press, Nottingham, UK.

Rodriguez, V. M. L., M. M. Calonge D, and D. Ordoñez Escudero. 2003. ELISA and HPLC determination of the occurrence of aflatoxin M1 in raw cow's milk. *Food Additives & Contaminants.* 20(3):276.

SAS Institute. 2009. SAS System for Windows. Ver. 9.1, SAS Institute Inc., Cary, NC.

Sadeghi, N. 2009. Incidence of aflatoxin M1 in human breast milk in Tehran, Iran. *Food Control.* 20(1):75.

Samson, A. R. 1992. Current taxonomic schemes of the genus *Aspergillus* and its teleomorphs. Pages 355-387 in *Aspergillus*. J. W. Bennett and M. A. Klich, eds. Butterworth-Heinemann, Stoneham, MA.

Sargent, K.A., J. O. K. Sheridan, and R. B. A. Carnaghan. 1961. Toxicity associated with certain samples of groundnuts. Page 4807 in *Nature*. Vol. 192.

Sarlmehmetoglu, B., O. Kuplulu, and T. Haluk Celik. 2004. Detection of aflatoxin M1 in cheese samples by ELISA. *Food Control* 15(1):45-49.

Scudamore, K. 2005. Principles and applications of mycotoxin analysis. Pages 157-186 in *The mycotoxin Blue Book*. D. Diaz, ed. Nottingham University Press, Nottingham, UK.

Selim, M. I., A. M. Juchems, and W. Popendorf. 1998. Assessing airborne aflatoxin B1 during on-farm grain handling activities. *Am. Indust. Hygiene Assoc. J.* 59(4):252-256.

Shah, V.P. (2007). The history of bioanalytical method validation and regulation: Evolution of a guidance document on bioanalytical methods validation. *The AAPS J.* 9 (1): E43-E47.

Sibanda, L., S. De Saeger, and C. Van Peteghem. 1999. Development of a portable field immunoassay for the detection of aflatoxin M1 in milk. *Int. J. Food Microbiol.* 48(3):203-209.

Sinha, K. K. 1998. Detoxification of mycotoxins and food safety. Pages 381-407 in *Mycotoxins in Agriculture and Food Safety*. K. K. Sinha and D. Bhatnagar, eds. Marcel Dekker, Inc., New York, NY.

Soha, S., M. T. Mazloumi, and M. Borji. 2006. Reduction of aflatoxin M1 residue in milk utilizing chemisorption compounds and its effects on quality of milk. *J. Arab Neonatology Forum*. Online: www.fmhs.uaeu.ac.ae/neonatal/iss006/pap05.asp. Accessed: 9 July 2010.

Stora, C., I. Dvorackova, and N. Ayraud. 1983. Aflatoxin and Reye's syndrome. *J. Med.* 14(1):47-54.

Stroka, J., and E. Anklam. 2002. New strategies for the screening and determination of aflatoxins and the detection of aflatoxin-producing moulds in food and feed. *Trac Trends in Analytical Chemistry*. 21(2):90-95.

Stroka, J., M. Petz, U. Joerissen, and E. Anklam. 1999. Investigation of various extractants for the analysis of aflatoxin B1 in different food and feed matrices. *Food Addit. Contam.* 16(8):331-338.

Turner Designs, 2008. Fluorescence Measurements: Chapter 1 Fluorescence Theory. http://www.turnerdesigns.com/t2/doc/appnotes/998_0050/0050_c1.html. Accessed Jan. 09, 2009.

Van Egmond, H. P. and S. Dragacci. 2000. Liquid chromatographic method for aflatoxin M1 in milk. Pages 59-69 in *Methods in Molecular Biology*. Vol. 157. M.W. Trucksess and A.E. Pohland, eds. Human Press Inc., Totowa, NY.

Van Egmond, H. P., and M. A. Jonker. 2004. Worldwide regulations on aflatoxins. The situation in 2002. *Toxin Reviews* 23(2):273 - 293.

Van Eijkeren, J. C. H., M. I. Bakker, and M. J. Zeilmaker. 2006. A simple steady-state model for carry-over of aflatoxins from feed to cow's milk. *Food Addit. Contam.* 23(8):833-838.

Vicam, L. P. 1999. AflaTest: Instructional Manual. [http://biotic.myweb.hinet.net/index.files/GN-MC9508-5%20\(AflaTest\).pdf](http://biotic.myweb.hinet.net/index.files/GN-MC9508-5%20(AflaTest).pdf), Accessed Jan. 26, 2009.

Wang, X. W., S. P. Hussain, T.-I. Huo, C.-G. Wu, M. Forques, L. J. Hofseth, C. Brechot, and C. C. Harris. 2002. Molecular pathogenesis of human hepatocellular carcinoma. *Toxicology* 181-182:43-47.

Weidenborner, M. 2001. *Encyclopedia of Food Mycotoxins*. Springer, Berlin, New York, London.

Whitlow, L. W. 2005. Molds and mycotoxins in feedstuffs - Prevention and treatment. Pages 123-142 in *Florida Ruminant Nutrition Symposium*, Univ. Florida, Gainesville.

Williams, J. H., T. D. Phillips, P. E. Jolly, J. K. Stiles, C. M. Jolly, and D. Aggarwal. 2004. Human aflatoxicosis in developing countries: A review of toxicology, exposure, potential health consequences, and interventions. *Am. J. Clin. Nutr.* 80(5):1106-1122.

Wilson, D. M., E. W. Sydenham, G. A. Lombaert, M. W. Truksess, D. Abramson, and G. A. Bennett. 1998. Mycotoxin analytical techniques. Pages 1441-1444 in *Mycotoxins in Agriculture and Food Safety*. K. K. Sinha and D. Bhatnagar, eds. Marcel Dekker, New York, NY.

Wu, F., Y. Liu, and D. Bhatnagar. 2008. Cost effectiveness of aflatoxin control methods: economic incentives. *Toxin Reviews* 27(3):203 - 225.

Yiannikouris, A., and J.-P. Jouany. 2002. Mycotoxins in feeds and their fate in animals: A review. *Animal Res.* 51(2):81-99.

Yin, Y., Y. Lei-yan., J. Jin-hua, and M. Zhong-hua. 2008. Biological control of aflatoxin contamination of crops. *J. Zhejiang Univ. Sci. B* 9(10):787-792.

Yousef, A. E., and E. H. Marth. 1989. Stability and degradation of aflatoxin M1. Pages 127-161 in *Mycotoxins in Dairy Products*. H. P. V. Egmond, ed. Elsevier, London, UK.