

A Simple and Efficient Approach to Quinazolinones under Mild Copper-Catalyzed Conditions**

Xiaowei Liu, Hua Fu,* Yuyang Jiang, and Yufen Zhao

Quinazolinone (**Q**) is a key core structure that occurs in natural products such as luotonin A from *Peganum nigellastrum*,^[1a] 2-methyl-4(3*H*)-quinazolinone from *Bacillus cereus*,^[1b] 2-(4-hydroxybutyl)quinazolin-4-one from *Dichroa febrifuga*,^[1c] and bouchardatine from *Bouchardatia neurococca* (Figure 1).^[1d] Quinazolinone derivatives are now




Figure 1. Quinazolinone (**Q**) and examples of related natural products.


known to have useful biological and medicinal activities; they can be used as hypnotic, sedative, analgesic, anticonvulsant, antitussive, antibacterial, antidiabetic, anti-inflammatory, and antitumor agents.^[2,3] Additionally, some therapeutic agents containing this core structure have been on the market or are in clinical trials for the treatment of cancer.^[4]

Although some methods for the synthesis of quinazolinone derivatives^[2,5,6] have been developed, they depend on the availability of the requisite *ortho*-amino- or -nitrobenzoic

acid derivatives. Some of the starting materials are also sometimes difficult to prepare. Recently, progress has been made on the copper-catalyzed Ullmann N-arylations,^[7] and they have been used to make N-heterocycles.^[8] Unfortunately, these methods are not useful for constructing some quinazolinone molecules because the reaction temperatures are too high. Therefore, it is desirable to develop milder copper-catalyzed coupling methods. Recently, Shafir and Buchwald^[9] and ourselves^[10] have developed copper-catalyzed N-arylations at room temperature, and the results showed that the efficiency of the copper-catalyzed coupling reactions was highly dependent on the involvement of suitable ligands. To the best of our knowledge, there is no example of constructing N-heterocycles under ligand-free copper catalysis at room temperature. Herein, we report a simple, practical, and efficient strategy for the synthesis of quinazolinone derivatives by using mild copper-catalyzed conditions in the absence of ligands or additives.

2-Bromobenzoic acid (**1a**) and acetamidine hydrochloride (**2a**) were chosen as the model substrates for the optimization of the reaction conditions, which include the catalyst, base, and solvent. As shown in Table 1, four copper catalysts were tested at room temperature by using two equivalents of Cs₂CO₃ as the base (relative to amount of **1a**) in DMF

Table 1: Copper-catalyzed coupling of 2-bromobenzoic acid with acetamidine hydrochloride: Optimization of the reaction conditions.^[a]



Entry	Catalyst	Base	Solvent	Yield [%] ^[b]
1	Cu	Cs ₂ CO ₃	DMF	19
2	CuSO ₄	Cs ₂ CO ₃	DMF	23
3	CuBr	Cs ₂ CO ₃	DMF	75
4	CuI	Cs ₂ CO ₃	DMF	81
5	CuI	Cs ₂ CO ₃	DMF	57 ^[c]
6	CuI	K ₂ CO ₃	DMF	74
7	CuI	K ₃ PO ₄	DMF	69
8	CuI	Cs ₂ CO ₃	toluene	trace
9	CuI	Cs ₂ CO ₃	dioxane	62
10	CuI	Cs ₂ CO ₃	DMSO	59
11	—	Cs ₂ CO ₃	DMF	trace ^[d]
12	CuI	Cs ₂ CO ₃	DMF	trace ^[e]

[a] Reaction conditions: 2-bromobenzoic acid (0.5 mmol), acetamidine hydrochloride (0.75 mmol), catalyst (0.1 mmol), base (1 mmol), solvent (3 mL) at room temperature (ca. 25 °C) under a nitrogen atmosphere. [b] Yield of isolated product. [c] Base (0.5 mmol). [d] No addition of catalyst. [e] Without nitrogen atmosphere. DMSO = dimethylsulfoxide; DMF = *N,N*-dimethylformamide.






[*] X. Liu, Prof. Dr. H. Fu, Prof. Dr. Y. Jiang, Prof. Dr. Y. Zhao
Key Laboratory of Bioorganic Phosphorus Chemistry and Chemical
Biology (Ministry of Education), Department of Chemistry
Tsinghua University, Beijing 100084 (China)
Fax: (+86) 10-6278-1695
E-mail: fuhua@mail.tsinghua.edu.cn

Prof. Dr. Y. Jiang
Key Laboratory of Chemical Biology (Guangdong Province)
Graduate School of Shenzhen, Tsinghua University
Shenzhen (China)

[**] Financial support was provided by the National Natural Science
Foundation of China (Grant No. 20672065), Chinese 863 Project
(Grant No. 2007AA02Z160), the Program for New Century Excellent
Talents in University (NCET-05-0062), and Changjiang Scholars and
Innovative Research Team in University (PCSIRT) (No. IRT0404) in
China and the Key Subject Foundation from the Beijing Department
of Education (XK100030514).

Supporting information for this article is available on the WWW
under <http://dx.doi.org/10.1002/ange.200804675>.

Table 2: (Continued)

Entry	1	2	Product	Yield [%] ^[b]
14	1d	2d		61
15		2a	3a	89
16	1e	2b	3b	90
17	1e	2d	3d	97
18	1e		3o	59 ^[c] (72 ^[d])
19	1e		3p	45 ^[c] (70 ^[d])
20		2b	3b	40 ^[c] (81 ^[d])
21	1f	2d	3d	43 ^[c] (85 ^[d])

[a] Reaction conditions: **1** (0.5 mmol), **2** (0.75 mmol), CuI (0.1 mmol), Cs₂CO₃ (1 mmol), DMF (3 mL) at room temperature (ca. 25 °C) under nitrogen atmosphere. [b] Yield of isolated product. [c] Reaction temperature of 40 °C. [d] Reaction temperature of 80 °C.

the previously reported copper-catalyzed N-arylations,^[7–11] and the result above demonstrates that an *ortho*-substituent effect is present during the N-arylation and is derived from the coordination of the carboxyl group (Scheme 1). The reaction of 2-bromo-5-chlorobenzoic acid (**1c**) with amidines took place at the C–Br bond *ortho* to the carboxyl group and not at the C–Cl bond, again demonstrating the presence of an *ortho*-substituent effect. The substituted 2-halobenzoic acids containing electron-rich groups showed slightly weaker reactivity than those containing electron-neutral or electron-deficient groups; for example, 6-bromo-1,3-benzodioxole-5-carboxylic acid (**1d**) provided lower yields (Table 2, entries 13 and 14). In general, amidines are good substrates, but the couplings of guanidines with 2-halobenzoic acids did not perform well at room temperature. The corresponding

quinazolinones were obtained in good yields when temperature was raised to 80 °C (Table 2, entries 18 and 19).

Quinazolinones can be readily transformed into the corresponding quinazolines, which have various biological and medicinal activities,^[11] so the present method provides a novel strategy for synthesis of a diverse array of quinazolinone and quinazoline derivatives.

Given that suitable *ortho* substituents promote Ullmann-type couplings,^[12] a mechanism for the formation of quinazolinones is proposed in Scheme 1. This proposal is based on the results discussed herein and on the other experimental evidence (see the Supporting Information for details). The coordination of substituted 2-haloben-


zoic acid to CuI forms **I** in the presence of the base (Cs₂CO₃). Oxidative addition of **I** and complexation of copper with the amidine provides **II**, which then undergoes reductive elimination to give the N-arylation product **III** and releases the copper catalyst. The coupling of the carboxyl and amino groups in **III** affords target product **3**.^[13]

In summary, we have developed a simple and highly efficient method for the synthesis of quinazolinone derivatives. The coupling reactions of 2-bromo- and iodobenzoic acid derivatives with amidines performed well at room temperature without the addition of a ligand or an additive. The target products were also obtained in higher yields from the nonactive substrates, such as 2-chlorobenzoic acid (**1f**) or guanidines, when the reaction temperature was raised to 80 °C. The present method is economical and practical, and the starting materials are readily available. These advantages, relative to previous methods, provide an opportunity for the construction of diverse and useful molecules within organic and medicinal chemistry.

Received: September 24, 2008

Published online: December 3, 2008

Keywords: copper · cyclization · guanidines · heterocycles · quinazolinone



Scheme 1. Proposed mechanism for the copper-catalyzed formation of quinazolinones.

- [1] a) Z.-Z. Ma, Y. Hano, T. Nomura, Y.-J. Chen, *Heterocycles* **1997**, 46, 541; b) S. Yoshida, T. Aoyagi, S. Harada, N. Matsuda, T. Ikeda, H. Naganawa, M. Hamada, T. Takeuchi, *J. Antibiot.* **1991**,

- 44, 111; c) Y. Deng, R. Xu, Y. Ye, *J. Chin. Pharm. Sci.* **2000**, 9, 116; d) C. Wattanapiromsakul, P. I. Forster, P. G. Waterman, *Phytochemistry* **2003**, 64, 609.
- [2] a) A. Witt, J. Bergman, *Curr. Org. Chem.* **2003**, 7, 659, and references therein; b) S. B. Mhaske, N. P. Argade, *Tetrahedron* **2006**, 62, 9787, and references therein.
- [3] a) D. J. Connolly, D. Cusack, T. P. O'Sullivan, P. J. Guiry, *Tetrahedron* **2005**, 61, 10153; b) Z. Ma, Y. Hano, T. Nomura, *Heterocycles* **2005**, 65, 2203.
- [4] For example, Raltitrexed (Tomudex, marketed for colorectal cancer), Ispinesib (phase II for solid tumors), and Tempostatin (phase II for bladder cancer).
- [5] a) A. Kamal, K. S. Reddy, B. R. Prasad, A. H. Babu, A. V. Ramana, *Tetrahedron Lett.* **2004**, 45, 6517; b) C. Larksarp, H. Alper, *J. Org. Chem.* **2000**, 65, 2773.
- [6] For selected examples, see: a) J.-F. Liu, P. Ye, B. Zhang, G. Bi, K. Sargent, L. Yu, D. Yohannes, C. M. Baldino, *J. Org. Chem.* **2005**, 70, 6339; b) J.-F. Liu, M. Kaselj, Y. Isome, P. Ye, K. Sargent, K. Sprague, D. Cherrak, C. J. Wilson, Y. Si, D. Yohannes, S.-C. Ng, *J. Comb. Chem.* **2006**, 8, 7; c) A. D. Roy, A. Subramanian, R. Roy, *J. Org. Chem.* **2006**, 71, 382; d) C. L. Yoo, J. C. Fetting, M. J. Kurth, *J. Org. Chem.* **2005**, 70, 6941.
- [7] For recent reviews on copper-catalyzed N-arylation, see: a) K. Kunz, U. Scholz, D. Ganzer, *Synlett* **2003**, 2428; b) S. V. Ley, A. W. Thomas, *Angew. Chem.* **2003**, 115, 5558; *Angew. Chem. Int. Ed.* **2003**, 42, 5400; c) I. P. Beletskaya, A. V. Cheprakov, *Coord. Chem. Rev.* **2004**, 248, 2337; d) G. Evano, N. Blanchard, M. Toumi, *Chem. Rev.* **2008**, 108, 3054, and references therein.
- [8] For recent studies on the syntheses of N-heterocycles by using Ullmann-type couplings, see: a) R. Martín, R. Rodríguez, S. L. Buchwald, *Angew. Chem.* **2006**, 118, 7237; *Angew. Chem. Int. Ed.* **2006**, 45, 7079; b) G. Evindar, R. A. Batey, *J. Org. Chem.* **2006**, 71, 1802; c) F. Bonnaterre, M. Bois-Choussy, J. Zhu, *Org. Lett.* **2006**, 8, 4351; d) B. Zou, Q. Yuan, D. Ma, *Angew. Chem.* **2007**, 119, 2652; *Angew. Chem. Int. Ed.* **2007**, 46, 2598; e) X. Yuan, X. Xu, X. Zhou, J. Yuan, L. Mai, Y. Li, *J. Org. Chem.* **2007**, 72, 1510; f) T. Yang, C. Lin, H. Fu, Y. Jiang, Y. Zhao, *Org. Lett.* **2005**, 7, 4781.
- [9] A. Shafir, S. L. Buchwald, *J. Am. Chem. Soc.* **2006**, 128, 8742.
- [10] a) D. Jiang, H. Fu, Y. Jiang, Y. Zhao, *J. Org. Chem.* **2007**, 72, 672; b) Q. Jiang, D. Jiang, Y. Jiang, H. Fu, Y. Zhao, *Synlett* **2007**, 1836.
- [11] L. Örfi, F. Wączek, J. Pató, I. Varga, B. Hegymegi-Barakonyi, R. A. Houghten, G. Kéri, *Curr. Med. Chem.* **2004**, 11, 2549, and references therein.
- [12] a) K. C. Nicolaou, C. N. C. Boddy, S. Natarajar, T.-Y. Yue, H. Li, S. Bräse, J. M. Ramanjulu, *J. Am. Chem. Soc.* **1997**, 119, 3421; b) J. Lindley, *Tetrahedron* **1984**, 40, 1433; c) A. V. Kalinin, J. F. Bower, P. Riebel, V. Snieckus, *J. Org. Chem.* **1999**, 64, 2986; d) Q. Cai, B. Zou, D. Ma, *Angew. Chem.* **2006**, 118, 1298; *Angew. Chem. Int. Ed.* **2006**, 45, 1276.
- [13] S. P. Flanagan, R. Goddard, P. J. Guiry, *Tetrahedron* **2005**, 61, 9808.